

the tissue is such that we were unable to have an equal age distribution. We also acknowledge that the three age groups are made up of subjects from different areas, and that these are not longitudinal data. Despite the above limitations, we consider the large number of subjects, whose deaths were not respiratory related, and the wide age range included provides a strong basis for establishing the time of appearance and pattern of thickening that occurs to the bronchial RBM, and its positive relationship with epithelial height, during normal development. These data indicate that future paediatric and adult biopsy studies assessing RBM thickness should include age-matched controls.

Funding The British Paediatric Respiratory Society, Asthma UK, The Wellcome Trust, UK. SS is funded by an Intermediate Clinical Fellowship from the Wellcome Trust. AJ is supported by a National Health and Medical Research Council of Australia Practitioner Fellowship.

Competing interests None.

Ethics approval This study was conducted with the approval of the local ethics committees at each centre from which tissue was obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

1. **Kirschvink N**, Reinhold P. Use of alternative animals as asthma models. *Curr Drug Targets* 2008;**9**:470–84.
2. **Evans MJ**, Fanucchi MV, Plopper CG, *et al*. Postnatal development of the lamina reticularis in primate airways. *Anat Rec (Hoboken)* 2010;**293**:947–54.
3. **Jeffery PK**. Remodeling in asthma and chronic obstructive lung disease. *Am J Respir Crit Care Med* 2001;**164**:S28–38.
4. **Payne DN**, Rogers AV, Adelroth E, *et al*. Early thickening of the reticular basement membrane in children with difficult asthma. *Am J Respir Crit Care Med* 2003;**167**:78–82.
5. **Barbato A**, Turato G, Baraldo S, *et al*. Airway inflammation in childhood asthma. *Am J Respir Crit Care Med* 2003;**168**:798–803.
6. **Saglani S**, Payne DN, Zhu J, *et al*. Early detection of airway wall remodeling and eosinophilic inflammation in preschool wheezers. *Am J Respir Crit Care Med* 2007;**176**:858–64.
7. **Holgate ST**, Davies DE, Lackie PM, *et al*. Epithelial–mesenchymal interactions in the pathogenesis of asthma. *J Allergy Clin Immunol* 2000;**105**:193–204.
8. **Hislop AA**, Haworth SG. Airway size and structure in the normal fetal and infant lung and the effect of premature delivery and artificial ventilation. *Am Rev Respir Dis* 1989;**140**:1717–26.
9. **Evans MJ**, Fanucchi MV, Van Winkle LS, *et al*. Fibroblast growth factor-2 during postnatal development of the tracheal basement membrane zone. *Am J Physiol Lung Cell Mol Physiol* 2002;**283**:L1263–70.
10. **Wilson JW**, Li X. The measurement of reticular basement membrane and submucosal collagen in the asthmatic airway. *Clin Exp Allergy* 1997;**27**:363–71.
11. **Jeffery PK**, Godfrey RW, Adelroth E, *et al*. Effects of treatment on airway inflammation and thickening of basement membrane reticular collagen in asthma. A quantitative light and electron microscopic study. *Am Rev Respir Dis* 1992;**145**:890–9.
12. **James AL**, Maxwell PS, Pearce-Pinto G, *et al*. The relationship of reticular basement membrane thickness to airway wall remodeling in asthma. *Am J Respir Crit Care Med* 2002;**166**:1590–5.
13. **James AL**, Pare PD, Hogg JC. The mechanics of airway narrowing in asthma. *Am Rev Respir Dis* 1989;**139**:242–6.
14. **Sullivan P**, Stephens D, Ansari T, *et al*. Variation in the measurements of basement membrane thickness and inflammatory cell number in bronchial biopsies. *Eur Respir J* 1998;**12**:811–15.
15. **Holgate ST**. Has the time come to rethink the pathogenesis of asthma? *Curr Opin Allergy Clin Immunol* 2010;**10**:48–53.
16. **Jeffery PK**. Embryology and growth of the normal lung. In: Brewis RAL, Gibson GJ, Geddes D, eds. *Respiratory Medicine*. London: Bailliere Tindall, 1990:3–20.
17. **Thurlbeck WM**. Postnatal human lung growth. *Thorax* 1982;**37**:564–71.
18. **Hislop AA**, Wigglesworth JS, Desai R, *et al*. The effects of preterm delivery and mechanical ventilation on human lung growth. *Early Hum Dev* 1987;**15**:147–64.
19. **Emery JL**, Mithal A. The number of alveoli in the terminal respiratory unit of man during late intrauterine life and childhood. *Arch Dis Child* 1960;**35**:544–7.
20. **Thurlbeck WM**, Angus GE. Growth and aging of the normal human lung. *Chest* 1975;**67**(2 Suppl):3S–6S.
21. **Ochs M**, Nyengaard JR, Jung A, *et al*. The number of alveoli in the human lung. *Am J Respir Crit Care Med* 2004;**169**:120–4.
22. **Shebani E**, Shahana S, Janson C, *et al*. Attachment of columnar airway epithelial cells in asthma. *Tissue Cell* 2005;**37**:145–52.
23. **Haworth SG**, Hislop AA. Lung development—the effects of chronic hypoxia. *Semin Neonatol* 2003;**8**:1–8.
24. **Saglani S**, Molyneux C, Gong H, *et al*. Ultrastructure of the reticular basement membrane in asthmatic adults, children and infants. *Eur Respir J* 2006;**28**:505–12.

Pulmonary puzzle

An unusual cause of chest pain

INTRODUCTION

A 19-year-old male patient with a history of asthma was in good health until he developed a sudden-onset, right side pleuritic chest discomfort and shortness of breath. At that time, he was

on a cruise to Puerto Rico. He returned home 1 week later and was evaluated in the emergency room at that time. On further history, he denied cough, fever, chills, night sweats, weight loss, sputum production, haemoptysis or other symptoms. He did not recall recent sick contacts or exposure to tuberculosis. He was born in the USA and this cruise was his only recent travel. He worked in a law firm and denied significant occupational exposures, any tobacco or drug use, allergies or significant family

Figure 1 (A) Chest radiograph showing significant mediastinal shift and volume loss on the right side, with prominent vascular markings only seen on the right side, (B) Chest CT scan showing right-sided volume loss with hypolucency, prominent right interstitial markings, subsegmental atelectasis, very small right pleural effusion and small right hilar and mediastinal adenopathy.





Figure 2 (A) Mucosal endobronchial biopsy: bronchial tissue with prominent submucosal capillaries and mild fibrosis. (B) Transbronchial biopsy: lung parenchyma with mild interstitial chronic inflammation and alveolar macrophages.



history of lung or cardiovascular disease. Tachycardia was the only abnormality on his physical examination, and initial laboratory testing were completely normal. His pulse oximetry measurement was 97% O₂ saturation on room air at rest. His ECG was consistent with sinus tachycardia, with a rate of 108/min and no ST–T wave changes. He was admitted to the hospital for further investigation of his chest pain.

His chest radiograph (figure 1A) and chest CT scan (angiogram protocol) (figure 1B) were abnormal but showed no evidence of pneumothorax or pulmonary embolism. A small unilateral pleural effusion was present but was too small to sample using bedside ultrasonography. All subsequent blood tests in the hospital including cardiac enzymes, immunoglobulins levels and vasculitis screen were all unremarkable. The chest x-ray and CT scan were remarkable for the predominance of right-sided

findings with relative sparing of the left side. This suggested the possibility of unilateral interstitial lung disease. Therefore, the patient underwent bronchoscopy to further narrow the differential diagnosis. Airway examination revealed prominent submucosal blood vessels with mucosal oedema and hyperaemia involving the surface of the right side airways with complete sparing of the left side airways. The biopsies revealed interstitial oedema and early fibrotic changes (figure 2A,B). On further review of the CT scan, an extraneous septum within the left atrium was noted and the diagnosis was made (figure 3).

QUESTION

What is the diagnosis?

See page 313 for the answer

Brian Casserly,¹ Michael K Atalay,² Athena Poppas,³ James R Klinger,⁴ Muhanned Abu-Hijleh⁴

¹Division of Pulmonary, Critical Care and Sleep Medicine, The Brown Alpert Medical School, Rhode Island, USA; ²Department of Diagnostic Imaging, Rhode Island Hospital, The Brown Alpert Medical School, Rhode Island, USA; ³Department of Cardiology, Rhode Island Hospital, The Brown Alpert Medical School, Rhode Island, USA; ⁴Division of Pulmonary, Critical Care and Sleep Medicine, Memorial Hospital of Rhode Island and Rhode Island Hospital, The Brown Alpert Medical School, Rhode Island, USA

Correspondence to Brian Casserly, Division of Pulmonary, Critical Care and Sleep Medicine, Memorial Hospital of Rhode Island, 111 Brewster Street, Pawtucket, RI 02860, USA; brian_casserly@brown.edu

Institution at which work was performed Rhode Island Hospital, Providence, RI 02903.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Published Online First 11 November 2010

Thorax 2011;66:285–286. doi:10.1136/thx.2010.141689



Figure 3 Chest CT scan that shows the septum in the left atrium.

7. **Hocking WG**, Hu P, Oken MM, *et al*. Lung cancer screening in the randomized Prostate, Lung, Colorectal, and Ovarian (PLCO) cancer screening trial. *J Natl Cancer Inst* 2010;**102**:722–31.
8. **Henschke CI**, Yankelevitz DF, Libby DM, *et al*. Survival of patients with stage I lung cancer detected on CT screening. *N Engl J Med* 2006;**355**:1763–71.
9. **Mulshine JL**, Sullivan DC. Lung cancer screening. *N Engl J Med* 2005;**352**:2714–20.
10. **Gray JA**, Patnick J, Blanks RG. Maximising benefit and minimising harm of screening. *BMJ* 2008;**336**:480–3.
11. **van den Bergh KA**, Essink-Bot ML, Borsboom GJ, *et al*. Short-term health-related quality of life consequences in a lung cancer CT screening trial (NELSON). *Br J Cancer* 2010;**102**:27–34.
12. **Whynes DK**. Could CT screening for lung cancer ever be cost effective in the United Kingdom. *Cost Eff Resour Alloc* 2008;**6**:5.
13. **Gohagan JK**, Marcus PM, Fagerstrom RM, *et al*. Final results of the Lung Screening Study, a randomized feasibility study of spiral CT versus chest X-ray screening for lung cancer. *Lung Cancer* 2005;**47**:9–15.
14. **Croswell JM**, Kramer BS, Kreimer AR, *et al*. Cumulative incidence of false-positive results in repeated, multimodal cancer screening. *Ann Fam Med* 2009;**7**:212–22.
15. <http://www.cancer.gov/newscenter/pressreleases/NLSTResultsRelease>.
16. **Aberle DR**, Berg CD, Black WC, *et al*; National Lung Screening Trial Research Team. The National Lung Screening Trial: overview and study design. *Radiology* 2011;**258**:243–53. <http://www.ncbi.nlm.nih.gov/pubmed/21045183>.
17. **Lopes Pegne A**, Picozzi G, Mascalchi M. Design, recruitment and baseline results of the ITALUNG trial for lung cancer screening with low-dose CT. *Lung Cancer* 2009;**64**:34–40.
18. **Infante M**, Cavuto S, Lutman F, *et al*. A randomized study of lung cancer screening with spiral computed tomography: three-year results from the DANTE trial. *Am J Respir Crit Care Med* 2009;**180**:445–53.
19. **Blanchon T**, Brechot JM, Grenier PA, *et al*. Baseline results of the Depiscan study: a French randomized pilot trial of lung cancer screening comparing low dose CT scan (LDCT) and chest X-ray (CXR). *Lung Cancer* 2007;**58**:50–8.
20. **Pedersen J**, Ashraf H, Dirksen A, *et al*. The Danish randomized lung cancer CT screening trial—overall design and results of the prevalence round. *J Thorac Oncol* 2009;**5**:608–14.
21. **Xu DM**, Gietema H, de Koning H, *et al*. Nodule management protocol of the NELSON randomised lung cancer screening trial. *Lung Cancer* 2006;**54**:177–84.
22. **Cassidy A**, Myles J, van-Tongeren M, *et al*. The LLP risk model: an individual risk prediction model for lung cancer. *Br J Cancer* 2008;**98**:270–6.
23. **Raji OY**, Duffy SW, Agbaje OF, *et al*. Predictive Accuracy and Clinical Utility of the Liverpool Lung Project Risk Model using Population-based Data from Europe and North America. Liverpool, UK: NCRI Conference, 2010.
24. **Yankelevitz DF**, Kostis WJ, Henschke CI, *et al*. Overdiagnosis in chest radiographic screening for lung carcinoma: frequency. *Cancer* 2003;**97**:1271–5.
25. **Lindell RM**, Hartman TE, Swensen SJ, *et al*. Five-year lung cancer screening experience: CT appearance, growth rate, location, and histologic features of 61 lung cancers. *Radiology* 2007;**242**:555–62.
26. **van Klaveren RJ**, Oudkerk M, Prokop M, *et al*. Management of lung nodules detected by volume CT scanning. *N Engl J Med* 2009;**361**:2221–9.
27. **Bach PB**, Kattan MW, Thornquist MD, *et al*. Variations in lung cancer risk among smokers. *J Natl Cancer Inst* 2003;**95**:470–8.
28. **van Klaveren RJ**, de Koning HJ, Mulshine J, *et al*. Lung cancer screening by spiral CT. What is the optimal target population for screening trials? *Lung Cancer* 2002;**38**:243–52.
29. **Spitz MR**, Hong WK, Amos CI, *et al*. A risk model for prediction of lung cancer. *J Natl Cancer Inst* 2007;**99**:715–26.
30. **Field JK**, Smith DL, Duffy S, *et al*. The Liverpool Lung Project research protocol. *Int J Oncol* 2005;**27**:1633–45.

Pulmonary puzzle

ANSWER

From the question on page 285

This is an unusual case of cor triatriatum sinistrum with secondary unilateral pulmonary venous hypertension and right lung hypoplasia. The interstitial changes in the parenchyma were secondary to a state of chronic congestion as a result of reduced interstitial fluid drainage due to elevated pressures in the right pulmonary vein. Our biopsy specimens highlight how chronic persistent interstitial oedema can result in chronic inflammation and the development of interstitial fibrosis.^{1,2} The enlarged mediastinal lymph nodes were probably hyperplastic and the gross findings on bronchoscopy reflected compensatory hypertrophy of the bronchial circulation draining into the right pulmonary vein.

Transoesophageal echocardiogram revealed a membrane in the left atrium (LA), preferentially obstructing the right pulmonary

venous inflow. The obstruction was incomplete, and high velocity flow across the membrane suggested elevated right pulmonary venous pressure of 25–30 mm Hg (figure 4A,B). This was confirmed by cardiac MRI which showed asymmetric pulmonary blood flow, with ~15–20 times more flow to the left lung than the right, with delay enhancement within the LA (figure 5). In our patient, despite the paucity of literature, surgical options were carefully considered. The patient underwent median sternotomy with surgical excision of the fibromuscular intra-atrial septum due to concerns about further progression of the pulmonary venous hypertension and the subsequent effects on the interstitium of his right lung. His postoperative course was unremarkable, and a 3-month follow-up cardiac MRI showed that the LA membrane can no longer be seen and there is no longer evidence of delayed enhancement within the LA (figure 6). Also, of note, there was a marked

Figure 4 (A) Echocardiogram: apical four-chamber view showing the septum (arrow) and extra atrial chamber (asterisk) (B) Echocardiogram: colour Doppler enhanced showing different flow velocities within the left atrium (arrow).




Figure 5 Preoperative cardiac MRI showing the left atrium (LA) membrane and the flow differential between the right and left pulmonary veins. Serial axial MRIs obtained during infusion of contrast demonstrate prompt enhancement and washout of the left inferior pulmonary vein (arrow), distal left atrium (dLA) and left ventricle (LV). The right inferior pulmonary vein (arrowhead) and proximal left atrium (pLA) show delayed enhancement. The LA membrane can be seen in image (b).





Figure 6 Serial axial MRIs obtained during infusion of contrast demonstrate gradual uniform enhancement of the left atrium (LA) and left ventricle (LV). The interatrial membrane has been resected. Note: the right pulmonary veins, while present were not seen in this slice location.



improvement in the flow differential between the right and left pulmonary veins on MRI.

DISCUSSION

To our knowledge, this is the first report in the literature of a patient with unilateral pulmonary venous hypertension and unilateral hypoplastic lung related to an atypical cor triatriatum sinistrum. Therefore, it is very important to distinguish this unusual case from other reported cases of classic cor triatriatum sinistrum. Typically, all of the pulmonary venous drainage enters

the proximal chamber and then crosses the septum through small fenestration(s) into the distal chamber (figure 7A). As a consequence, pulmonary venous hypertension (bilateral) develops and results in recurrent pulmonary oedema. This clinical syndrome resembles (and is frequently misdiagnosed as) congenital mitral stenosis.³ Surgical resection in these circumstances is the optimal therapeutic option.⁴

In our patient, the intra-atrial septum separated the right from the left pulmonary veins, creating a small chamber in the left atrium that had the right pulmonary venous return (figure 7B). A pin-hole fenestration connected this chamber with the supramitral left atrium that received left pulmonary venous return. This created significant isolated pulmonary venous congestion and pulmonary venous hypertension in the right lung with significant difference in pulmonary blood flow between the right and left lungs as demonstrated at cardiac MRI.

This case highlights the need to consider a cardiac aetiology in patients with prominent unilateral interstitial lung markings on chest x-ray or CT scan, and how echocardiography and cardiac MRI are valuable tools in the diagnosis and follow-up of such conditions. Surgical resection of the membrane appears to be the optimal treatment option for all age groups.

Thorax 2011;**66**:313–314. doi:10.1136/thx.2010.141689




Figure 7 (A) Classic cor triatriatum sinistrum: illustration of the pulmonary veins entering the proximal chamber with the incoming blood escaping through a small fenestration(s) that communicates with the distal left atrium. (B) Unusual cor triatriatum sinistrum (our patient): the extra chamber created by the intra-atrial septum separated the right from the left pulmonary veins by a membrane with a pin-hole fenestration, creating significant pulmonary venous congestion and hypertension in the right lung.

REFERENCES

1. Drake RE, Doursout MF. Pulmonary edema and elevated left atrial pressure: four hours and beyond. *News Physiol Sci* 2002;**17**:223–6.
2. Parker S, Weiss F. The nature and significance of the structural changes in the lungs in mitral stenosis. *Am J Pathol* 1936;**22**:573–98.
3. Marin-Garcia J, Tandon R, Lucas RV Jr, et al. Cor triatriatum: study of 20 cases. *Am J Cardiol* 1975;**35**:59–66.
4. Alphonso N, Norgaard MA, Newcomb A, et al. Cor triatriatum: presentation, diagnosis and long-term surgical results. *Ann Thorac Surg* 2005;**80**:1666–71.