

ORIGINAL ARTICLE

Mesenchymal stem cells enhance recovery and repair following ventilator-induced lung injury in the rat

Gerard F Curley,^{1,2} Mairead Hayes,^{1,2} Bilal Ansari,^{1,2} Georgina Shaw,³ Aideen Ryan,³ Frank Barry,³ Timothy O'Brien,³ Daniel O'Toole,^{2,3} John G Laffey^{1,2,3}

See Editorial, p 475

► Additional materials are published online only. To view these files please visit the journal online (<http://thorax.bmj.com>).

¹Department of Anaesthesia, School of Medicine, Clinical Sciences Institute, Galway, Ireland

²Lung Biology Group, National Centre for Biomedical Engineering Sciences, Galway, Ireland

³Regenerative Medicine Institute, National University of Ireland, Galway, Ireland

Correspondence to

Professor John G Laffey, Anaesthesia, School of Medicine, Clinical Sciences Institute, National University of Ireland, Galway, Ireland; john.laffey@nuigalway.ie

Data from this paper were presented at the American Thoracic Society Annual Scientific Meeting, Denver, Colorado, May 2011.

Received 6 September 2011
Accepted 27 October 2011
Published Online First
21 November 2011

ABSTRACT

Background Bone-marrow derived mesenchymal stem cells (MSCs) reduce the severity of evolving acute lung injury (ALI), but their ability to repair the injured lung is not clear. A study was undertaken to determine the potential for MSCs to enhance repair after ventilator-induced lung injury (VILI) and elucidate the mechanisms underlying these effects.

Methods Anaesthetised rats underwent injurious ventilation which produced severe ALI. Following recovery, they were given an intravenous injection of MSCs (2×10^6 cells) or vehicle immediately and a second dose 24 h later. The extent of recovery following VILI was assessed after 48 h. Subsequent experiments examined the potential for non-stem cells and for the MSC secretome to enhance VILI repair. The contribution of specific MSC-secreted mediators was then examined in a wound healing model.

Results MSC therapy enhanced repair following VILI. MSCs enhanced restoration of systemic oxygenation and lung compliance, reduced total lung water, decreased lung inflammation and histological lung injury and restored lung structure. They attenuated alveolar tumour necrosis factor α concentrations while increasing concentrations of interleukin 10. These effects were not seen with non-stem cells (ie, rat fibroblasts). MSC-secreted products also enhanced lung repair and attenuated the inflammatory response following VILI. The beneficial effect of the MSC secretome on repair of pulmonary epithelial wounds was attenuated by prior depletion of keratinocyte growth factor.

Conclusion MSC therapy enhances lung repair following VILI via a paracrine mechanism that may be keratinocyte growth factor-dependent.

INTRODUCTION

Acute lung injury (ALI) and acute respiratory distress syndrome (ARDS) are devastating diseases with a mortality of up to 40%, and for which there are no treatments.¹ Mortality from ALI/ARDS has fallen² as a result of advances in supportive care for sepsis, trauma and pneumonia and the demonstration that mechanical ventilation, while necessary for survival, has the capacity to cause significant harm.³ The importance of ventilator-induced lung injury (VILI) is underscored by the fact that ventilation strategies that reduce lung stretch save lives.³ The mechanisms whereby ventilation contributes to lung injury are increasingly well understood.⁴ However, more recent attempts to adjust ventilation strategies to further

Key messages

What is the key question?

- Are bone marrow mesenchymal stem cells (MSCs) capable of repairing the lung following severe injury by injurious mechanical ventilation?

What is the bottom line?

- Bone marrow MSCs enhance repair in rat lungs following injurious ventilation and encourage restitution in pulmonary epithelial wounds by a paracrine mechanism involving secretion of keratinocyte growth factor.
- These effects were not seen with non-stem cells, were mediated via a paracrine mechanism, involved restoration of the pulmonary epithelium and were dependent in part on MSC secretion of keratinocyte growth factor.

Why read on?

- This study highlights the capacity of MSCs to repair the lung following ventilation-induced acute lung injury.

reduce harm have met with limited success.⁵ Even with contemporary low stretch strategies, it appears difficult to avoid regional areas of high lung stretch.⁶ In addition, low stretch strategies may worsen atelectasis,⁷ which can also cause harm.

An alternative approach is to develop strategies that enhance lung repair following VILI. Mesenchymal stem cells (MSCs) are fibroblast-like cells that can be isolated from bone marrow and are characterised by their ability to self-renew and undergo differentiation into mesenchymal lineage cell types including bone, cartilage, adipose tissue, muscle and tendon.⁸ MSCs have shown promise in a number of preclinical ALI/ARDS studies^{9–10} and appear to exert immunomodulatory, anti-inflammatory and regenerative effects.⁹

The potential for MSCs to augment wound healing and repair after stretch-induced lung injury is not known. However, several factors suggest that they play a key role in the reparative response following injury,¹¹ and raise the possibility that transplanted MSCs may enhance restoration of organ function following injury. MSCs secrete growth factors and cytokines, which may play an important role in tissue regeneration and repair.¹² They can be found in increased numbers under




Figure 1 Flow diagram indicating timelines for experimental interventions. MSCs, mesenchymal stem cells; PBS, phosphate-buffered saline; VILI, ventilator-induced lung injury.

stress conditions, and both endogenous¹¹ and transplanted¹³ MSCs appear to home to sites of injury. MSCs enhance tissue regeneration and wound repair in preclinical disease models and in clinical studies.¹⁴ In recent clinical studies, MSCs accelerated healing of cutaneous wounds and increased epithelialisation and angiogenesis.¹⁵

Given these issues, we hypothesised that MSCs would enhance functional and structural recovery after VILI. We further hypothesised that this effect would be mediated, at least in part, via MSC-secreted soluble factors.

MATERIALS AND METHODS

All work was approved by the Animal Ethics Committee, National University of Ireland, Galway, and conducted under licence from the Department of Health, Ireland. A full description of the methods is given in the online supplement.

MSC harvest and cell culture

Rat MSCs (rMSCs) were isolated from adult male Sprague-Dawley rats. Following aspiration, the bone marrow was plated into tissue culture flasks. Adherent cells were grown until 80%

confluent and then trypsinised and culture expanded. Fibroblasts, used as control cells, were obtained from the dermis of adult Sprague-Dawley rats after digestion in 0.25% trypsin.

Human MSCs (hMSCs) were aspirated from the iliac crests of healthy human volunteers and cultured until 80% confluent, when they were harvested and expanded. Primary human lung fibroblasts and A549 lung adenocarcinoma cells were obtained from American Type Culture Collection (ATCC). MSCs were characterised according to international guidelines (see figures A and B in online supplement).¹⁶

Conditioned medium

Rat and human MSCs and fibroblasts (2×10^6) were washed and cultured without serum for 24 h. The cells were again washed and the subsequent serum-free medium for the next 24 h was used as the conditioned medium (CM). For in vivo experiments, 15 ml of this medium was concentrated using a 3000 Da centrifugal concentrating filter (Amicon, Billerica, Massachusetts, USA) to give 500 μ l.


Rodent ventilator-induced lung injury protocol

The timeline for these experiments is depicted in figure 1. As previously described, adult male Sprague Dawley rats were anaesthetised, orotracheally intubated and subjected to injurious mechanical ventilation (inspiratory pressure 35 cm H₂O, respiratory rate 18/min and positive end expiratory pressure 0 cm H₂O).¹⁷ When respiratory static compliance had decreased by 50%, the animals were recovered and entered into the treatment protocol.¹⁷

Assessment of injury and repair

At 48 h following VILI induction, animals were re-anaesthetised and arterial blood gases and static inflation lung compliance

Figure 2 Mesenchymal stem cell (MSC) therapy enhances resolution following ventilation-induced lung injury. MSC therapy (A) decreased alveolar–arterial oxygen gradient ($p=0.04$), (B) increased static lung compliance ($p=0.003$), (C) reduced lung wet to dry weight ratios ($p=0.01$) and (D) decreased bronchoalveolar lavage (BAL) protein concentrations ($p=0.03$) 48 h after induction of severe stretch-induced lung injury compared with vehicle ($n=8$ animals per group). * $p<0.05$ vs vehicle.



measurements were performed.¹⁸ Post-mortem, bronchoalveolar lavage (BAL) differential cell counts, protein concentration and cytokine levels were determined. Wet to dry lung weight ratios were measured and histological lung damage was assessed.

In vivo experimental series

In the first series, after VILI the animals were randomly allocated to receive an intravenous injection of either 2×10^6 allogeneic rMSCs in 500 μ l phosphate buffered saline (PBS) or 500 μ l PBS alone. Twenty-four hours later a second injection of rMSCs or vehicle was administered (figure 1). In the second series, after VILI and at 24 h, animals received injections of 2×10^6 rMSCs, 500 μ l PBS, 2×10^6 rat dermal fibroblasts or 500 μ l rMSC-CM.

Wound repair experiments

An in vitro model of alveolar epithelial repair was used, as used in prior studies.¹⁹ Single wounds were made in confluent A549 monolayers in 24 well plates with a 1000 μ l pipette tip.¹⁹ The wounds were exposed to different conditions according to group allocation. In the first experiment the wounds were incubated in MEM- α medium, human fibroblast CM, human MSC (hMSC)-CM or co-cultured with hMSCs. In the second experiment, hMSC-CM was incubated with monoclonal antibodies to inactivate keratinocyte growth factor (KGF), hepatocyte growth factor (HGF) and transforming growth factor β (TGF- β) (Abcam, Cambridge, UK), respectively. Wounds were exposed to hMSC-CM with and without antibodies to each candidate mediator. At 48 h the extent of epithelial restitution was determined (Photoshop v8.0, Adobe Systems, San Jose, California, USA).

Statistical analysis

Data were analysed using Sigma Stat (SYSTAT Software, Richmond, California, USA). The distribution of all data was

tested for normality using Kolmogorov–Smirnov tests. Data were analysed by one-way ANOVA followed by the Dunnett test, with the vehicle group as the control group in each analysis. Comparisons between two groups were made using unpaired two-tailed Student *t* tests or a Mann–Whitney *U* test as appropriate. Underlying model assumptions were deemed appropriate on the basis of suitable residual plots. A two-tailed *p* value of <0.05 was considered significant.

RESULTS

In the first experimental series, 18 animals were entered into the study. Two animals died post-injury prior to randomisation, leaving eight animals per group. There were no differences between the groups at baseline with regard to animal weight and duration of injurious ventilation required to induce injury.

MSCs restore lung function following VILI

The administration of rMSCs enhanced recovery following VILI. rMSCs improved arterial oxygenation, as measured by the alveolar–arterial oxygen gradient (figure 2A), and enhanced restoration of static compliance compared with vehicle controls (figure 2B). rMSCs enhanced the restoration of lung microvascular permeability, as evidenced by a decrease in lung wet to dry weight ratios (figure 2C) and a decrease in alveolar fluid protein concentrations (figure 2D).

MSCs modulate inflammation following VILI

rMSCs reduced both total BAL cell count and BAL neutrophil counts (figure 3A). Alveolar concentrations of tumour necrosis factor α (TNF α) (figure 3B) but not interleukin 6 (IL-6) (figure 3C) were decreased following rMSC treatment. In contrast,

Figure 3 Mesenchymal stem cell (MSC) therapy modulates the inflammatory response to ventilation-induced lung injury. MSC therapy (A) decreased bronchoalveolar lavage (BAL) neutrophil counts ($p=0.045$), (B) decreased BAL tumour necrosis factor α (TNF α) concentrations ($p=0.04$), (C) did not alter BAL interleukin 6 (IL-6) concentrations ($p=0.42$) and (D) increased BAL interleukin 10 (IL-10) concentrations ($p=0.02$) 48 h after induction of severe stretch-induced lung injury compared with vehicle ($n=8$ animals per group). * $p<0.05$ vs vehicle.





Figure 4 Mesenchymal stem cell (MSC) therapy enhances the resolution of structural lung injury following ventilator-induced lung injury. MSC therapy enhanced resolution of histological injury as evidenced by (A) decreased alveolar lung tissue and (B) increased alveolar airspace fraction. Representative photomicrographs of lung from (C) a vehicle-treated animal and (D) a MSC-treated animal show greater resolution of lung injury with MSCs at 48 h (n=8 animals per group). Scale bar 200 μ m. *p<0.05 vs vehicle.



alveolar concentrations of the anti-inflammatory cytokine IL-10 were increased in response to rMSC therapy (figure 3D).


MSCs restore lung structure following VILI

rMSCs decreased alveolar thickening, as evidenced by reduced alveolar tissue volume fraction, and increased recovery of airspace volume, as evidenced by increased alveolar airspace volume fraction (figures 4A,B). Representative histological sections of lung demonstrate the greater degree of resolution of injury and alveolar infiltrates in rMSC-treated animals (figure 4C,D).

MSCs enhance repair via a paracrine mechanism

In the second series, 40 animals (N=10 per group) underwent induction of severe VILI. There were no differences between the groups at baseline. Both rMSCs and rMSC-CM enhanced repair following VILI. In contrast, fibroblasts did not appear to exert beneficial effects. rMSCs and rMSC-CM enhanced the recovery of static lung compliance following VILI (figure 5A) and significantly reduced alveolar total cell and neutrophil infiltration (figure 5B), alveolar TNF α concentrations (figure 5C) and alveolar IL-6 concentrations (figure 5D). rMSCs and rMSC-CM significantly

Figure 5 Mesenchymal stem cells (MSCs) exert benefits via a paracrine mechanism. Both MSCs and MSC conditioned medium (MSC-CM) (A) enhanced recovery of static lung compliance, (B) decreased bronchoalveolar lavage (BAL) neutrophil counts, (C) decreased BAL tumour necrosis factor α (TNF α) concentrations and (D) reduced alveolar interleukin 6 (IL-6) concentrations 48 h after injury compared with animals that received fibroblasts or fibroblast conditioned medium (Fibroblast-CM) (n=10 animals per group). *p<0.05 vs fibroblast CM and fibroblast groups.



increased serum, but not alveolar, IL-10 concentrations compared with animals treated with fibroblasts (data not shown).

MSCs enhance pulmonary epithelial wound repair via a KGF-dependent mechanism

Human MSC-CM increased the rate of wound closure in alveolar epithelial A549 monolayers subjected to scratch injury in comparison with fibroblast CM and fresh medium controls (figure 6A). The rate of wound closure seen with hMSC-CM was similar to that seen in hMSC co-cultures (figure 6A). In subsequent studies, prior incubation of hMSC-CM with antibodies to neutralise KGF attenuated its beneficial effects on wound repair (figure 6B). In contrast, incubation of hMSC-CM with antibodies to neutralise HGF and TGF- β did not alter wound repair (figure 6B). Measurement of KGF concentrations in the MSC and fibroblast CM showed that MSCs produced substantially greater amounts of KGF (510 ± 90 vs 68 ± 13 pg/ml, $p < 0.0001$).

DISCUSSION

MSCs have several properties that make them attractive as a potential treatment for ALI/ARDS. MSCs avoid allo-recognition,

home to sites of injury and suppress inflammation as well as immune responses.^{9 10 20} Preclinical studies have shown that MSC therapy can attenuate endotoxin-induced ALI when given during the injury phase in both rodents¹⁰ and in the perfused human lung.²¹ Bone marrow-derived MSCs have also demonstrated benefit in preclinical sepsis models.⁹

In these studies we have shown that MSCs enhance the recovery of lung function and structure following VILI. Specifically, rMSCs reduced the decrement in systemic oxygenation, restored static lung compliance, and enhanced alveolar fluid clearance following VILI, as evidenced by reduced BAL protein concentrations, reduced lung wet to dry weight ratios and histological evidence of reduced alveolar tissue oedema. These findings support previous findings that rMSCs enhance alveolar epithelial fluid transport in the endotoxin-injured ex vivo human lung via a KGF-dependent mechanism.²¹ The demonstration that rMSCs enhanced the recovery of lung function and structure following VILI is central to determining its therapeutic potential. Many promising therapies have demonstrated benefit in preclinical models when used prophylactically but have been less successful when used following injury.

MSCs decreased the inflammatory response following VILI. Specifically, rMSCs reduced alveolar inflammatory cell counts, particularly alveolar neutrophils. It is not clear whether rMSCs reduced alveolar neutrophil infiltration or enhanced neutrophil clearance. These findings support those of previous studies, which suggest that MSCs modulate the activity of T cells, B cells, dendritic cells, monocytes and macrophages.^{20 22–24} Raffaghello *et al*²⁵ recently reported that MSCs inhibited apoptosis and the oxidative burst of resting and activated neutrophils while preserving their phagocytic and chemotactic functions.

In our studies, MSCs modulated the cytokine response, decreasing alveolar TNF α and IL-6 concentrations. rMSC therapy increased alveolar IL-10 concentrations, a potentially important finding which has previously been reported.¹⁰ IL-10 is a cytokine secreted predominantly by monocytes that decreases the expression of Th1 cytokines, MHC class II antigens and co-stimulatory molecules on macrophages. IL-10 also inhibits the rolling, adhesion and transepithelial migration of neutrophils.²⁶ Nemeth *et al* described a pivotal role for MSC-induced macrophage secretion of IL-10 in mediating the protective effects of MSCs in a murine caecal ligation and puncture sepsis model.⁹ Nemeth *et al* further showed that MSCs induced macrophage IL-10 secretion via secretion of prostaglandin E2.

MSCs enhanced repair following VILI via a paracrine mechanism, with rMSC-CM as effective as rMSCs themselves. To further dissect the mechanisms of action of MSCs, we focused on the potential for MSC secretion of KGF, TGF- β and HGF to contribute to repair. We found that depletion of KGF—but not TGF- β or HGF—from the MSC medium attenuated pulmonary epithelial wound healing. Given the importance of epithelial wound healing to restoration of lung structure following VILI and the previous demonstration that MSCs enhance alveolar fluid clearance via a KGF-dependent mechanism,²¹ these findings suggest that KGF may be central to the reparative effects of MSCs in VILI.

The dose regimen used for these studies—namely, intravenous 2×10^6 rMSCs following recovery from VILI followed by a second dose 24 h later—was safe and effective with no adverse effects related to rMSC administration. However, other dosage regimens and other delivery routes such as the intratracheal^{10 21} or intraperitoneal routes²⁷ may be equally effective in restoring lung function following VILI. Dermal fibroblasts were used as




Figure 6 Mesenchymal stem cells (MSCs) enhance pulmonary epithelial wound closure via a KGF-dependent mechanism. (A) In pulmonary epithelial layers, both MSCs and MSC conditioned medium (MSC-CM) enhanced wound closure. (B) Incubation of MSC-CM with antibodies to neutralise KGF, but not HGF or TGF- β , attenuated its effect on wound healing ($n=12$ per group). KGF, keratinocyte growth factor; HGF, hepatocyte growth factor; TGF- β , transforming growth factor β . *Significantly different from fibroblast CM or control group. †Significantly different from MSC-CM groups.

a control non-stem cell because of their similarity to MSCs.²⁸ Additional studies are needed to determine the optimal dosage regimen, including the minimally effective MSC doses and the best administration route.

There are a number of limitations to these studies. First, while we provide data to suggest that KGF secreted by MSCs contributes to repair following VILI, our data do not preclude a contribution by other mediators to the mechanisms by which MSCs enhance lung repair. Second, our in vivo studies were carried out in a rodent model and caution must be exercised in extrapolating to the clinical situation. However, our in vitro studies were carried out in human cell lines and used human MSCs. When taken together with other studies examining the effects of MSCs during the injury phase of ALI/ARDS, these findings strongly suggest that MSCs may have therapeutic potential for ALI/ARDS. Third, we did not provide baseline data on these animals to allow the reader to assess the magnitude of effects of injurious ventilation on the parameters measured. However, the effect of the high stretch ventilation strategy used in this model has been well characterised in a previous publication from our group.¹⁷ Furthermore, we do not provide data for the effects of MSCs on protectively ventilated or unventilated lungs. However, it would be expected that any effects on these animals would be limited. Lastly, we did not study the effects of KGF supplemented or depleted MSC-CM in our in vivo studies. Inclusion of these groups might have provided important additional insights.

In conclusion, we have shown that bone marrow-derived MSCs enhance recovery after VILI when administered into the systemic circulation in the rat. The mechanism for this effect may be due to the secretion of paracrine soluble factors such as KGF by the MSCs themselves. MSC therapy may represent an innovative approach for treatment of VILI and ARDS.

Funding The European Research Council (grant number ERC-2007-StG 207777) and Health Research Board, Dublin, Ireland (grant number RP/2008/193) provided unrestricted funds for these studies. Dr Curley was supported by a fellowship from Molecular Medicine Ireland (HEA PRTL Cycle A).

Competing interests None.











Patient consent Not needed.




Contributors JGL, GFC, TO'B, DOT and FB designed the study. GFC and MH performed the experiments. BA carried out the cytokine assays. GFC, AR and GS harvested and characterised the mesenchymal stem cells. JGL analysed the data. GFC, DOT and JGL drafted the manuscript and are guarantors of the paper.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

1. Rubenfeld GD, Herridge MS. Epidemiology and outcomes of acute lung injury. *Chest* 2007;**131**:554–62.
2. Zamboni M, Vincent JL. Mortality rates for patients with acute lung injury/ARDS have decreased over time. *Chest* 2008;**133**:1120–7.
3. ARDSNet. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. The Acute Respiratory Distress Syndrome Network. *N Engl J Med* 2000;**342**:1301–8.
4. Vlahakis NE, Hubmayr RD. Cellular stress failure in ventilator-injured lungs. *Am J Respir Crit Care Med* 2005;**171**:1328–42.
5. Meade MO, Cook DJ, Guyatt GH, et al. Ventilation strategy using low tidal volumes, recruitment maneuvers, and high positive end-expiratory pressure for acute lung injury and acute respiratory distress syndrome: a randomized controlled trial. *JAMA* 2008;**299**:637–45.
6. Terragni PP, Rosboch G, Tealdi A, et al. Tidal hyperinflation during low tidal volume ventilation in acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2007;**175**:160–6.
7. Fanelli V, Mascia L, Puntorieri V, et al. Pulmonary atelectasis during low stretch ventilation: "open lung" versus "lung rest" strategy. *Crit Care Med* 2009;**37**:1046–53.
8. Prockop DJ. Marrow stromal cells as stem cells for nonhematopoietic tissues. *Science* 1997;**276**:71–4.
9. Nemeth K, Leelahavanichkul A, Yuen PS, et al. Bone marrow stromal cells attenuate sepsis via prostaglandin E(2)-dependent reprogramming of host macrophages to increase their interleukin-10 production. *Nat Med* 2009;**15**:42–9.
10. Gupta N, Su X, Popov B, et al. Intrapulmonary delivery of bone marrow-derived mesenchymal stem cells improves survival and attenuates endotoxin-induced acute lung injury in mice. *J Immunol* 2007;**179**:1855–63.
11. Rochefort GY, Delorme B, Lopez A, et al. Multipotential mesenchymal stem cells are mobilized into peripheral blood by hypoxia. *Stem Cells* 2006;**24**:2202–8.
12. Prockop DJ. Repair of tissues by adult stem/progenitor cells (MSCs): controversies, myths, and changing paradigms. *Mol Ther* 2009;**17**:939–46.
13. Barhash IM, Chouraqui P, Baron J, et al. Systemic delivery of bone marrow-derived mesenchymal stem cells to the infarcted myocardium: feasibility, cell migration, and body distribution. *Circulation* 2003;**108**:863–8.
14. Yoshikawa T, Mitsuno H, Nonaka I, et al. Wound therapy by marrow mesenchymal cell transplantation. *Plast Reconstr Surg* 2008;**121**:860–77.
15. Wu Y, Chen L, Scott PG, et al. Mesenchymal stem cells enhance wound healing through differentiation and angiogenesis. *Stem Cells* 2007;**25**:2648–59.
16. Dominici M, Le Blanc K, Mueller I, et al. Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement. *Cytotherapy* 2006;**8**:315–17.
17. Curley GF, Contreras M, Higgins B, et al. Evolution of the Inflammatory and Fibroproliferative Responses during Resolution and Repair Following Ventilator-induced Lung Injury in the Rat. *Anesthesiology* 2011;**115**:1022–32.
18. Higgins BD, Costello J, Contreras M, et al. Differential effects of buffered hypercapnia versus hypercapnic acidosis on shock and lung injury induced by systemic sepsis. *Anesthesiology* 2009;**111**:1317–26.
19. O'Toole D, Hassett P, Contreras M, et al. Hypercapnic acidosis attenuates pulmonary epithelial wound repair by an NF-kappaB dependent mechanism. *Thorax* 2009;**64**:976–82.
20. Beyth S, Borovsky Z, Mevorach D, et al. Human mesenchymal stem cells alter antigen-presenting cell maturation and induce T-cell unresponsiveness. *Blood* 2005;**105**:2214–19.
21. Lee JW, Fang X, Gupta N, et al. Allogeneic human mesenchymal stem cells for treatment of E. coli endotoxin-induced acute lung injury in the ex vivo perfused human lung. *Proc Natl Acad Sci U S A* 2009;**106**:16357–62.
22. Aggarwal S, Pittenger MF. Human mesenchymal stem cells modulate allogeneic immune cell responses. *Blood* 2005;**105**:1815–22.
23. Glennie S, Soeiro I, Dyson PJ, et al. Bone marrow mesenchymal stem cells induce division arrest anergy of activated T cells. *Blood* 2005;**105**:2821–7.
24. Corcione A, Benvenuto F, Ferretti E, et al. Human mesenchymal stem cells modulate B-cell functions. *Blood* 2006;**107**:367–72.
25. Raffaghello L, Bianchi G, Bertolotto M, et al. Human mesenchymal stem cells inhibit neutrophil apoptosis: a model for neutrophil preservation in the bone marrow niche. *Stem Cells* 2008;**26**:151–62.
26. Ajuebor MN, Das AM, Virag L, et al. Role of resident peritoneal macrophages and mast cells in chemokine production and neutrophil migration in acute inflammation: evidence for an inhibitory loop involving endogenous IL-10. *J Immunol* 1999;**162**:1685–91.
27. Danchuk S, Ylostalo JH, Hossain F, et al. Human multipotent stromal cells attenuate lipopolysaccharide-induced acute lung injury in mice via secretion of tumor necrosis factor-alpha-induced protein 6. *Stem Cell Res Ther* 2011;**2**:27.
28. Haniffa MA, Collin MP, Buckley CD, et al. Mesenchymal stem cells: the fibroblasts' new clothes? *Haematologica* 2009;**94**:258–63.

CD29**CD90****CD44H****CD73****CD45RA****CD71****CD80****MHC-1****MHC-II****CD106**

A**B****C**

Online Data Supplement

Title: Mesenchymal Stem Cells enhance recovery and repair following Ventilator Induced Lung Injury **in the Rat**

Authors: Dr. Gerard Curley, MB, FCARCSI ^{1,2},
Dr Mairead Hayes, MB, FCARCSI^{1,2},
Dr Bilal Ansari, MB, FCARCSI^{1,2},
Ms Georgina Shaw ³,
Dr Aideen Ryan, PhD³,
Prof Frank Barry, PhD ³,
Prof Tim O'Brien, MD, PhD ³
Dr. Daniel O' Toole, PhD ^{1,2},
Prof. John G. Laffey, MD, FCARCSI ^{1,2,3}.

Institution: ¹Department of Anaesthesia, Clinical Sciences Institute; ²Lung Biology Group, National Centre for Biomedical Engineering Sciences; and ³Regenerative Medicine Institute, National University of Ireland, Galway, IRELAND.

Contact information: Prof John G Laffey, Anaesthesia, School of Medicine, Clinical Sciences Institute, National University of Ireland, Galway, Ireland.

E-mail: john.laffey@nuigalway.ie

Phone: 353 – 91 – 544908.

Fax: 353 – 91 – 544908.

Abbreviated Title: MSC's enhance repair following VILI

MATERIALS AND METHODS

All work was approved by the Animal Ethics Committee of the National University of Ireland, Galway and conducted under license from the Department of Health, Ireland. Specific-pathogen-free adult male Sprague Dawley rats (Charles River Laboratories, Kent, UK) weighing between 350–450g were used in all experiments.

Rodent MSC Isolation and Culture

Rodent MSCs (rMSCs) were isolated from rat femora and tibiae under sterile conditions as previously described (1) with additional modifications. Briefly, male Sprague Dawley rats (8-12 weeks old) were euthanized by inhalation of CO₂. Incisions were made on both lower limbs to expose the tibiae and femora. Both bones were removed from the hind limbs and placed in ice cold sterile Tyrode's solution (Sigma, St. Louis, MO). The marrow was then flushed into a dish containing rMSC complete culture medium (MEM- α Media (Gibco, Paisley, UK), F12-Ham Media (Gibco), 10% foetal bovine serum (PAA, Somerset, UK), 1% antibiotic/antimycotic (Sigma) and dispersed into a cell suspension. After centrifugation and filtration through a 100 μ m nylon mesh, a cell count was performed and the cells were transferred to a 175 cm² flasks containing 30 mls of rMSC complete medium, at a density of 9×10^5 cells/cm². On day 3 of culture in an atmosphere of 5% CO₂/90% humidity at 37°C, medium and non-adherent cells were removed and fresh medium was added to each flask. Cells were ready for subculture (usually after 16-17 days) when colonies began to exhibit a compact appearance and multi-layered growth or when the loosely formed colonies began to merge into a monolayer (<90% of confluence).

Thereafter, cells were ready to be passaged after 6/7 days culture, at 80% confluence. For passage, media was aspirated off and cells were washed with sterile PBS to remove any remaining serum. 8mls 0.25% trypsin/EDTA solution was added to the cells, which were incubated for 5 minutes at 37° C. The enzymatic reaction was stopped by adding the same volume of rMSC media to cells. Cells were centrifuged at 400g for 5 min. Media was aspirated off the cell pellet which was resuspended in 1ml and a haemocytometer count was undertaken. Cells were expanded to passage 4, whereupon they were used for experiments. (*Figure A, Online Repository.*)

Human MSC Isolation and Culture

Human MSCs were aspirated under sterile conditions from the iliac crests of healthy human volunteers. The obtained marrow was filtered with a 70 µm cell strainer (Falcon, USA) before centrifuging at 400 g for 10 min. Cell pellets were resuspended in media consisting MEM-α (Gibco), supplemented with 10% Fetal Bovine Serum (FBS) (Gibco, USA) and 1% antibiotics (streptomycin and penicillin) (Gibco, USA), and cultured in 175 cm² flasks at 37 °C in a humidified atmosphere containing 5% CO₂. At day 4, the cultures were washed with PBS to remove the non-adherent cells and further expanded until >80% confluence, when they were harvested and expanded in 175 cm² flasks. After subculture, these cells were designated as passage 1.

Characterization of MSCs

In accordance with the position statement for the minimal criteria to define an MSC (2), cells were labeled with monoclonal antibodies against CD31, CD34,

CD44, CD45, CD54, CD73, and CD90 (Santa Cruz Biotechnology, Santa Cruz, CA) and analyzed with a FACScan (Becton Dickinson, Franklin Lakes, NJ) and CellQuest software as described (3).

Osteogenic differentiation was induced by culturing rMSCs or hMSCs for up to 4 weeks in rMSC or hMSC Complete Medium supplemented with dexamethasone, ascorbic acid and β -glycerophosphate as previously described (4). To observe calcium deposition, cultures were stained with Alizarin Red stain (Sigma, St Louis, MA). To induce adipogenic differentiation, rMSCs were cultured for up to 4 weeks in medium supplemented with dexamethasone and insulin; adipocytes were discerned by staining with Oil Red O (Sigma). Chondrogenic capacity was assessed by addition of TGF-beta (Invitrogen) and staining with Toluidine blue (*Figure B panels a to c, Online Repository*).

Fibroblast isolation and Culture

Adult male Sprague Dawley rats were euthanized by CO₂ inhalation. The ventral surface of the rat was shaved and sprayed with 70% ethanol. Skin and subcutaneous tissue was removed and placed into 70% ethanol for 30 seconds. Fat and subcutaneous tissue was removed and the skin strips were placed in 0.25% trypsin (Sigma) overnight. The epidermis was then peeled from the dermal layer, and the dermal layer was placed on a scored 6 well plate (Sarstedt, Wexford, Ireland) in F-12/MEM- α medium supplemented with fetal calf serum (10%) and penicillin/streptomycin (1%).

Primary human lung fibroblasts were obtained from American Type Culture Collection (ATCC).

Conditioned Medium

Allogeneic human or rodent MSC, or human or rodent fibroblasts (2×10^6) were washed with PBS and cultured without serum for 24 h. The cells were again washed and the medium was then replaced, and the subsequent medium without serum for the next 24 h was used as the conditioned medium (CM). All conditioned medium was sterile filtered through a 22 μ m filter to remove cellular debris. For the *in vivo* experiments 15 mls of this medium was concentrated using a 3000 kDa centrifugal concentrating filter (Amicon, Billerica, MA, USA) to give 500 μ L.

Rodent Ventilator Induced Injury Protocol

Anesthesia was induced with intraperitoneal ketamine 80 mg.kg⁻¹ (Ketalar, Pfizer, Cork, Ireland) and xylazine 8 mg.kg⁻¹ (Xylapan, Vétquinol, Dublin, Ireland). After confirmation of depth of anesthesia by paw clamp, intravenous access was obtained via tail vein, laryngoscopy was performed and the animals were intubated with a size 14G intravenous catheter (BD Insyte®, Becton Dickinson Ltd, Oxford, UK). The lungs were ventilated using a small animal ventilator (CWE SAR 830 AP, CWE Inc, Pennsylvania, USA). Anesthesia was maintained with repeated boli of Saffan® (alfaxadone 0.9% and alfadadolone acetate 0.3%; Schering Plough, Welwyn Garden City, UK) and muscle relaxation was achieved with cis-atracurium besylate 0.5mg.kg⁻¹ (GlaxoSmithKline, Dublin, Ireland). The animals were then subjected to a high stretch mechanical ventilation protocol (FiO₂ 0.3, inspiratory pressure 35 cmH₂O, respiratory rate 18 min⁻¹, and positive end-expiratory pressure 0 cmH₂O). When static compliance

had decreased by 50%, high stretch ventilation was discontinued and animals were extubated, allowed to regain consciousness, and entered into the treatment protocol.

Assessment of Injury and Repair

At 48 hours following VILI induction, animals were re-anesthetized. A tracheostomy was inserted and carotid arterial access established (22G, BD Insyte), and the lungs were mechanically ventilated at a respiratory rate of 80 min⁻¹, tidal volume 6 ml.kg⁻¹ and positive end-expiratory pressure 2 cmH₂O as previously described (5-7). Intra-arterial blood pressure, peak airway pressures and rectal temperature were recorded continuously. Static inflation lung compliance measurements were performed as previously described (8, 9). After 20 minutes, the inspired gas was altered to a FiO₂ of 1.0 for 15 min, and a final arterial blood sample was taken. Heparin (400 IU.kg⁻¹, CP Pharmaceuticals, Wrexham, U.K.) was then administered intravenously, and animals were killed by exsanguination.

Immediately post-mortem, the heart-lung block was dissected and bronchoalveolar lavage (BAL) collection was performed as previously described (10, 11). BAL differential cell counts were performed. Protein concentration was determined using a Micro BCA™ Protein assay kit (Pierce, Rockford, IL, USA).(12) BAL IL-1β, IL-6, TNF-α and IL-10 concentrations were determined using quantitative sandwich enzyme-linked immunosorbent assays (R and D Systems, Abingdon, UK) (13). The detection limit of these assays was 62.5 pg/ml for the IL-6 assay and 31.2 pg/ml for TNF-α and IL-10 assays.

Wet:dry lung weight ratios were determined using the lowest lobe of the right lung as previously described (14). The left lung was isolated and fixed for morphometric examination, and the extent of histologic lung damage was determined using quantitative stereological techniques as previously described (10, 14).

Wound Repair Experiments

A549 cells were purchased from the European collection of cell cultures (Porton Down, UK) as cryopreserved 90-passage culture and used at passages 91-95. All cells were seeded at a density of 5×10^5 cells per ml on plastic 24 well plates (Corning Ltd, New York, US) at 37°C in a humidified incubator saturated with a gas mixture containing 5% CO₂ in air. Once the cells had grown to confluence, single wounds were made in each well, by scraping off cells with a 1000µL pipette tip, as previously described (18). Wells were washed with PBS and exposed to different conditions as per group allocation. 48 hours later, the monolayers were fixed with 4% paraformaldehyde in PBS (w/v), and stained with hematoxylin and eosin. The extent of epithelial restitution was determined by imaging each plate on a flatbed scanner and assessing the area of each wound using edge-finding software (Photoshop v8.0, Adobe Systems Inc, San Jose, California).

Experimental series

Series 1: Determination of potential for MSCs to enhance repair post VILI

Following recovery after induction of VILI, animals were randomly allocated to receive a tail vein injection of either: (1) 2×10^6 allogeneic rat MSCs suspended in 500 μ L PBS; or (2) 500 μ L PBS alone, and returned to cages. 24 hours later they were re-anesthetized, a second injection of MSCs or vehicle was administered intravenously, and the animals returned to their cages. The extent of repair following VILI assessed at 48 hours.

Series 2: Determination of the mechanism by which MSCs enhance repair

Following induction of VILI, animals were randomly allocated to receive a tail vein injection of: (1) 2×10^6 MSCs suspended in 500 μ L PBS; (2) 500 μ L PBS; (3) 2×10^6 rat dermal fibroblasts suspended in PBS; or (4) 500 μ L of MSC conditioned medium, and were returned to cages. 24 hours later they received a second injection of MSCs, vehicle, fibroblasts or medium. The extent of recovery and repair following VILI was assessed at 48 hours.

Series 3: Effects of MSCs and MSC-CM on epithelial wound repair

Pulmonary epithelial (A549) monolayers were subjected to scratch wound injury, incubated in: (i) MEM- α medium, (ii) human fibroblast conditioned medium, (iii) human MSC conditioned medium, or (iv) co-cultured with MSCs (Corning HTS transwell, pore size 0.4 μ m, Corning, NY, USA). In the case of co-cultures, MSCs were seeded at 1×10^3 cells/cm² in the inserts and maintained in human MSC medium for 3 days prior to co-culture, which allowed the mesenchymal stem cells to reach 70–80% confluence. Mesenchymal stem cell containing inserts were washed and then added to the A549 wells and flooded with fresh serum free medium. Similarly, as controls, human fibroblasts were

seeded at 1×10^3 cells in a co-culture insert and maintained in MSC medium for 3 days prior to co-culture.

Wound closure was assessed at 48 hours.

Series 4: Determination of the contribution of MSC secreted growth factors to wound repair.

MSC conditioned medium was incubated with specific monoclonal antibodies to inactivate keratinocyte growth factor (KGF), hepatocyte growth factor (HGF) and transforming growth factor-beta (TGF- β) (Abcam) respectively. The concentration of antibody used was sufficient to ensure effective blockade based on the expected amounts of the specific growth factor in conditioned medium. In the case of KGF, used 2 μ g/ml of this antibody given that 0.56-0.85 μ g/ml of this antibody is required to yield 50% inhibition of the biological activity of 25 ng/ml hKGF. Measurement of the concentration of KGF in the MSC and fibroblast conditioned medium, using quantitative sandwich enzyme-linked immunosorbent assays (R and D Systems, Abingdon, UK) confirmed that the concentration of neutralizing antibody used was sufficient. A549 wounds were exposed to MSC conditioned medium with and without antibodies to each candidate mediator, and the extent of wound closure assessed at 48 hours.

Statistical Analysis

The distribution of all data was tested for normality using Kolmogorov-Smirnov tests. Results are expressed as mean (\pm SD) for normally distributed data, and as median (interquartile range, IQR) where non-normally distributed. Data were

analyzed by one-way ANOVA followed by Dunnett's test, with the vehicle group as the control group in each analysis. Comparisons between 2 groups were made using unpaired, two-tailed Student's t tests or using a Mann-Whitney U test as appropriate. Underlying model assumptions were deemed appropriate on the basis of suitable residual plots. A two-tailed p value of <0.05 was considered significant.

FIGURE LEGENDS

Figure A: Characterization of surface markers on rat MSCs

The rat MSCs used in these studies were CD29, CD90, CD44H, CD73 positive and CD45RA, CD71, CD80, MHCI, MHCII, CD106 low or negative. Shown are FACS histograms of Sprague Dawley MSCs (passage 3) stained with antibodies against surface markers as indicated (colored) or with appropriate isotype controls (gray). Each colored line indicates replicates.

Figure B: Differentiation of rat MSCs

Passage 2 bone marrow (BM)-derived adherent cells (Figure B-a), after culture in differentiation medium, were stained for alizarin red (Osteogenic staining, Figure B-b), oil red O (Adipocyte staining, Figure B-c) and safranin O (Chondrocyte staining, data not shown). All experiments were performed in triplicate.

References

1. Meirelles Lda S, Nardi NB. Murine marrow-derived mesenchymal stem cell: Isolation, in vitro expansion, and characterization. *Br J Haematol* 2003;123:702-711.
2. Dominici M, Le Blanc K, Mueller I, et al. Minimal criteria for defining multipotent mesenchymal stromal cells. The international society for cellular therapy position statement. *Cytotherapy* 2006;8:315-317.
3. Rochefort GY, Delorme B, Lopez A, et al. Multipotential mesenchymal stem cells are mobilized into peripheral blood by hypoxia. *Stem Cells* 2006;24:2202-2208.
4. Phinney DG, Kopen G, Isaacson RL, et al. Plastic adherent stromal cells from the bone marrow of commonly used strains of inbred mice: Variations in yield, growth, and differentiation. *J Cell Biochem* 1999;72:570-585.
5. Curley G, Contreras M, Higgins B, O'Kane C, McAuley DF, O' Toole D, Laffey JG. Evolution of the inflammatory and fibroproliferative responses during resolution and repair following ventilator induced lung injury in the rat. *Anesthesiology* 2011.
6. Higgins BD, Costello J, Contreras M, et al. Differential effects of buffered hypercapnia versus hypercapnic acidosis on shock and lung injury induced by systemic sepsis. *Anesthesiology* 2009;111:1317-1326.
7. O'Croinin DF, Nichol AD, Hopkins N, et al. Sustained hypercapnic acidosis during pulmonary infection increases bacterial load and worsens lung injury. *Crit Care Med* 2008;36:2128-2135.

8. Costello J, Higgins B, Contreras M, et al. Hypercapnic acidosis attenuates shock and lung injury in early and prolonged systemic sepsis. *Crit Care Med* 2009;37:2412-2420.
9. Higgins BD, Costello J, Contreras M, et al. Differential effects of buffered hypercapnia versus hypercapnic acidosis on shock and lung injury induced by systemic sepsis. *Anesthesiology* 2009;111:1317-1326.
10. O'Croinin DF, Hopkins NO, Moore MM, et al. Hypercapnic acidosis does not modulate the severity of bacterial pneumonia-induced lung injury. *Crit Care Med* 2005;33:2606-2612.
11. O'Croinin DF, Nichol AD, Hopkins N, et al. Sustained hypercapnic acidosis during pulmonary infection increases bacterial load and worsens lung injury*. *Crit Care Med* 2008;36:2128-2135.
12. Smith PK, Krohn RI, Hermanson GT, et al. Measurement of protein using bicinchoninic acid. *Analytical biochemistry* 1985;150:76-85.
13. Adamson IY, Bakowska J. Relationship of keratinocyte growth factor and hepatocyte growth factor levels in rat lung lavage fluid to epithelial cell regeneration after bleomycin. *Am J Pathol* 1999;155:949-954.
14. Laffey JG, Honan D, Hopkins N, et al. Hypercapnic acidosis attenuates endotoxin-induced acute lung injury. *Am J Respir Crit Care Med* 2004;169:46-56.