

The early origins of COPD in severe asthma: the one thing that leads to another or the two things that come together?

Joerg Mattes,^{1,2} Peter G Gibson²

Chronic obstructive pulmonary disease (COPD) is diagnosed during middle to late adult life, when significant airflow limitation has already developed by the time the disease is recognised. What if the origins of COPD occurred much earlier? If this were true, it would have major implications for detection (screening) and prevention of COPD. Convincing evidence is hard to obtain, because it requires exceptionally long follow-up, such as half a life span, a period that is often longer than the average career of a researcher! Williams and McNichol established such a longitudinal asthma study from a 1957 birth cohort, and now Tai *et al*¹ evaluate for an association between childhood asthma and adult COPD. They followed-up a cohort of 6–7-year-old children over more than four decades, of whom 197 survivors completed lung function testing and questionnaires at age 50 years. The most important result reported by the authors was that children with more severe asthma were at increased risk (OR 32; 95% CI, 3 to 269) of developing adult COPD. Specifically, of those who had more severe asthma in childhood, 44% (15/34) were diagnosed with adult COPD and 41% (14/34) with adult asthma, while 15% (5/34) showed asthma remission at age 50 years. By contrast, adult COPD affected only 8% (13/163) of those who had intermittent or no childhood asthma. Thus, adults who suffered from more severe asthma in childhood have more commonly COPD.

Several other results reported here are of interest. First, previous or current tobacco smoking status did not predict adult COPD in this cohort, and forced expiratory volume in 1 s/forced vital capacity (FEV1/FVC) ratios were not different

between smokers and non-smokers in the COPD group. To put this unexpected result into perspective, it is important to consider the study design. A random sampling strategy of a birth cohort encompassing 30 000 children was used to 'enrich' The Melbourne Asthma Cohort with children who have asthma, and to increase—by means of sampling a further 21 000 children—the number of those with severe asthma.^{2,3} Considering the relatively small size of this cohort along with the difficulty to ascertain smoking behaviours at multiple time points, it is conceivable that the study was underpowered to detect tobacco smoke-related COPD cases and lung function decline. Smoke-related COPD is quite common but affects only a minority of smokers. Instead, the Melbourne Asthma Cohort was sufficiently powered to detect cases with 'severe asthma-related COPD'. This is less common but appears to affect quite a large proportion of adults with a history of more severe asthma in childhood.

Second, no accelerated decline in FEV1 or FEV1/FVC was observed in subjects with COPD as compared to those with asthma, or neither disease. This is an unexpected finding that could be explained by a lack of study power for this interaction, insufficient tobacco dose in the population, disease heterogeneity, or positive effects of treatment for asthma. Recent studies show heterogeneity in the degree of lung function decline in treated COPD. It is likely that the widespread use of inhaled corticosteroids in adult asthma has modified the loss of lung function. Alternatively, the mechanism for the loss of lung function may not be accelerated decline of lung function in adult life, but rather, incomplete airway growth during childhood. There is support for this in the Busselton Health study,⁴ which showed incomplete attainment of lung function in those with childhood asthma. Potential mechanisms include repeated insults from severe asthma due to repeated attacks of eosinophilic inflammation⁵ or bronchoconstriction⁶ that promote airway remodelling, or as yet

undefined mechanisms that modify the developing airway.

Third, severe childhood asthma did not increase the risk for having asthma at age 50 years (defined as having asthma symptoms and taking asthma medications for the past 3 years) compared to intermittent childhood asthma. Specifically less than half the children with asthma had current asthma in adulthood, irrespective of severity. This is comparable to asthma remission rates in previous studies, that is, ~20–30% during a 10-year period.^{7,8} Therefore, the majority of adults with asthma represent the minority of subjects who have not outgrown childhood-onset asthma. What the results from Tai *et al* add is that most cases with severe childhood-onset asthma who had an apparent asthma remission have, in fact, progressed to adult COPD defined according to the Global Initiative for Chronic Obstructive Lung Disease (GOLD) criteria (post-bronchodilator FEV1/FVC ratio less than 0.7).⁹ However, whether a complete asthma remission has occurred would require more assessments over time, because gaps in asthma activity, even for several years, are very common.¹⁰

There is broad consensus that classical asthma and tobacco smoke-related COPD are clinically, immunologically and histopathologically distinct. However, an overlap between asthma and COPD in some patients is well recognised.¹¹ Overlap is associated with more symptoms, exacerbations, hospitalisations and lower health-related quality of life, possibly independent of lung function and emphysema.^{12–14} Thus, spirometry has a limited efficacy in detecting this increased symptom burden and asthma overlap. Biomarkers, such as neutrophil gelatinase-associated lipocalin (NGAL) in sputum may be more suitable, because NGAL is higher in some COPD patients as compared with normal individuals,¹⁵ and also differentiates those with overlap syndrome from other COPD phenotypes.¹⁶ This may be relevant to the findings from Tai *et al* because enhanced neutrophilic or mixed granulocytic inflammation is also found more commonly in severe asthma.¹⁷ Potentially severe asthma is associated with COPD-asthma overlap, which would require further molecular, cellular and physiological studies.

Irrespective of this 'one thing leads to another' versus 'two things come together' conundrum, the Melbourne Asthma Cohort study demonstrates unequivocally that impaired lung function is established in childhood and tracks throughout life. This is supported by data from the unselected Tuscon birth cohort identifying impaired airway function shortly after birth as a risk

¹Experimental & Translational Respiratory Medicine, University of Newcastle, Newcastle, New South Wales, Australia; ²Hunter Medical Research Institute, University of Newcastle, Newcastle, New South Wales, Australia

Correspondence to Professor Joerg Mattes, HMRI, Newcastle Children's Hospital, Locked Bag 1, Newcastle, NSW 2305, Australia; joerg.mattes@newcastle.edu.au

factor for airflow obstruction in young adults.¹⁸ These studies support the hypothesis that there is a window of opportunity for developing interventions that promote respiratory health to prevent incomplete lung growth in early life—and consequently, COPD. Reducing tobacco smoke exposure is the obvious one as parental and active smoking act synergistically to affect early lung function deficits in young adulthood.¹⁹ Notably, several COPD genes modify responses to smoke exposure and are involved in lung growth and development in early life.²⁰ Reversing the global tobacco epidemic remains the great challenge as more than 90% of the world's population is not protected by comprehensive smoke-free policy.²¹ Exposure to respiratory viruses in early life is also very common and—if the result is a more severe lower respiratory tract infection (LRTI)—constitutes an independent risk factor for lung function abnormalities in adulthood.²² Maternal smoking and asthma are risk factors for virus-induced LRTIs in infancy.²³ Interestingly, less asthma exacerbations during pregnancy through treatment adjustments guided by inflammometry dramatically reduced LRTI in their offspring,^{24 25} but the effects on lung growth have yet to be determined. The study from Tai *et al* provides another valuable and carefully executed piece of research evidence to strengthen the hypothesis that developing successful interventions to reduce adverse events in utero and in early life will result in healthier lung growth and less chronic lung disease in adulthood.

Contributors JM and PGG wrote the editorial.

Competing interests None.

Provenance and peer review Not commissioned; internally peer reviewed.



CrossMark

To cite Mattes J, Gibson PG. *Thorax* 2014;**69**:789–790.

Published Online First 9 May 2014



► <http://dx.doi.org/10.1136/thoraxjnl-2013-204815>

Thorax 2014;**69**:789–790.
doi:10.1136/thoraxjnl-2014-205401

REFERENCES

- Tai A, Tran H, Roberts M, *et al*. The association between childhood asthma and adult chronic obstructive pulmonary disease. *Thorax* 2014;**69**: 805–10.
- Williams H, McNicol KN. Prevalence, natural history, and relationship of wheezy bronchitis and asthma in children. An epidemiological study. *Br Med J* 1969;**4**:321–5.
- Martin AJ, McLennan LA, Landau LI, *et al*. The natural history of childhood asthma to adult life. *Br Med J* 1980;**280**:1397–400.
- James AL, Palmer LJ, Kicic E, *et al*. Decline in lung function in the Busselton Health Study: the effects of asthma and cigarette smoking. *Am J Respir Crit Care Med* 2005;**171**:109–14.
- Phipps S, Benyahia F, Ou TT, *et al*. Acute allergen-induced airway remodeling in atopic asthma. *Am J Respir Cell Mol Biol* 2004;**31**:626–32.
- Grainge CL, Lau LC, Ward JA, *et al*. Effect of bronchoconstriction on airway remodeling in asthma. *N Engl J Med* 2011;**364**:2006–15.
- Holm M, Omenaas E, Gislason T, *et al*. Remission of asthma: a prospective longitudinal study from northern Europe (RHINE study). *Eur Respir J* 2007;**30**:62–5.
- Cazzoletti L, Corsico AG, Albicini F, *et al*. The course of asthma in young adults: a population-based nine-year follow-up on asthma remission and control. *PLoS ONE* 2014;**9**:e86956.
- Celli BR, MacNee W. Standards for the diagnosis and treatment of patients with COPD: a summary of the ATS/ERS position paper. *Eur Respir J* 2004;**23**: 932–46.
- Gershon A, Guan J, Victor JC, *et al*. The course of asthma activity: a population study. *J Allergy Clin Immunol* 2012;**129**:679–86.
- Marsh SE, Travers J, Weatherall M, *et al*. Proportional classifications of COPD phenotypes. *Thorax* 2008;**63**:761–7.
- Menezes AM, Montes de Oca M, Perez-Padilla R, *et al*. Increased risk of exacerbation and hospitalization in subjects with an overlap phenotype: COPD-asthma. *Chest* 2014;**145**: 297–304.
- Rhee CK, Yoon HK, Yoo KH, *et al*. Medical utilization and cost in patients with overlap syndrome of chronic obstructive pulmonary disease and asthma. *Copd* 2014;**11**:163–70.
- Kauppi P, Kupiainen H, Lindqvist A, *et al*. Overlap syndrome of asthma and COPD predicts low quality of life. *J Asthma* 2011;**48**:279–85.
- Keatings VM, Barnes PJ. Granulocyte activation markers in induced sputum: comparison between chronic obstructive pulmonary disease, asthma, and normal subjects. *Am J Respir Crit Care Med* 1997;**155**:449–53.
- Iwamoto H, Gao J, Koskela J, *et al*. Differences in plasma and sputum biomarkers between COPD and COPD-asthma overlap. *Eur Respir J* 2014;**43**:421–9.
- Moore WC, Hastie AT, Li X, *et al*. Sputum neutrophil counts are associated with more severe asthma phenotypes using cluster analysis. *J Allergy Clin Immunol* Published Online First: 8 Dec 2013. doi: 10.1016/j.jaci.2013.10.011
- Stern DA, Morgan WJ, Wright AL, *et al*. *Lancet*. 2007;**370**:758–64.
- Guerra S, Stern DA, Zhou M, *et al*. Combined effects of parental and active smoking on early lung function deficits: a prospective study from birth to age 26 years. *Thorax* 2013;**68**:1021–8.
- Kerkhof M, Boezen HM, Granell R, *et al*. Transient early wheeze and lung function in early childhood associated with chronic obstructive pulmonary disease genes. *J Allergy Clin Immunol* 2014;**133**: 68–76 e1–4.
- World Health Organization. *Research for International Tobacco Control. WHO report on the global tobacco epidemic*. Geneva: World Health Organization, 2008.
- Korppi M, Piippo-Savolainen E, Korhonen K, *et al*. *Pediatr Pulmonol* 2004;**38**:155–60.
- Carroll KN, Gebretsadik T, Griffin MR, *et al*. Maternal asthma and maternal smoking are associated with increased risk of bronchiolitis during infancy. *Pediatrics* 2007;**119**:1104–12.
- Powell H, Murphy VE, Taylor DR, *et al*. Management of asthma in pregnancy guided by measurement of fraction of exhaled nitric oxide: a double-blind, randomised controlled trial. *Lancet* 2011;**378**:983–90.
- Mattes J, Murphy VE, Powell H, *et al*. Prenatal origins of bronchiolitis: protective effect of optimised asthma management during pregnancy. *Thorax* 2014;**69**:383–4.