ORIGINAL ARTICLE

Aberrant expression and activity of histone deacetylases in sporadic idiopathic pulmonary fibrosis

Martina Korfei,^{1,2} Sylwia Skwarna,^{1,2} Ingrid Henneke,^{1,2} BreAnne MacKenzie,^{1,2} Oleksiy Klymenko,^{1,2} Shigeki Saito,³ Clemens Ruppert,^{1,2,4} Daniel von der Beck,^{1,2} Poornima Mahavadi,^{1,2} Walter Klepetko,^{5,11} Saverio Bellusci,^{1,2,4} Bruno Crestani,^{6,11} Soni Savai Pullamsetti,^{1,2,7} Ludger Fink,^{2,4,8} Werner Seeger,^{1,2,4} Oliver Holger Krämer,⁹ Andreas Guenther^{1,2,4,10,11}

ABSTRACT Background Activation and differentiation of

fibroblasts into contractile protein-expressing

myofibroblasts and their acquired apoptosis-resistant

of idiopathic pulmonary fibrosis (IPF), a fatal disease

characterised by distorted pulmonary structure and

excessive extracellular matrix (ECM) deposition. The

remain incompletely understood. We investigated the

possible implication of aberrant overexpression and

sporadic IPF (n=26) and non-diseased control lungs

IPF fibroblasts were treated with HDAC inhibitors

(HDACi) LBH589 or valproic acid (VPA).

significantly elevated in IPF lungs. Using

(n=16) for expression of class I and II HDACs. Primary

Results Compared to control lungs, protein levels of

class I (HDAC1, HDAC2, HDAC3, HDAC8) and class II

HDACs (HDAC4, HDAC 5, HDAC 7, HDAC 9) were

immunohistochemistry, strong induction of nearly all HDAC enzymes was observed in mvofibroblasts of

fibroblast foci and in abnormal bronchiolar basal cells at

sites of aberrant re-epithelialisation in IPF lungs, but not

in controls. Treatment of primary IPF fibroblasts with the

proliferation and cell survival, as well as in suppression

pan-HDACi LBH589 resulted in significantly reduced

expression of genes associated with ECM synthesis,

of HDAC7, and was paralleled by induction of

endoplasmic reticulum stress and apoptosis. The profibrotic and apoptosis-resistant phenotype of IPF fibroblasts was also partly attenuated by the class I

Conclusions Aberrant overexpression of HDACs in basal cells of IPF lungs may contribute to the

generation and apoptosis resistance of IPF fibroblasts are

Therefore, pan-HDAC inhibition by LBH589 may present

bronchiolisation process in this disease. Similarly,

mediated by enhanced activity of HDAC enzymes.

a novel therapeutic option for patients with IPF.

activity of histone deacetylases (HDACs) in IPF.

molecular mechanisms underlying these processes in IPF

Methods We analysed lung tissues from patients with

phenotype are critical factors towards the development

► Additional material is published online only. To view please visit the journal online (http://dx.doi.org/10.1136/ thoraxjnl-2014-206411).

For numbered affiliations see end of article.

Correspondence to

Professor Andreas Guenther, Department of Internal Medicine II, Justus-Liebig-University Giessen, Klinikstrasse 36, Giessen D-35392, Germany; Andreas.Guenther@ innere.med.uni-giessen.de

Received 6 October 2014 Revised 10 August 2015 Accepted 15 August 2015 Published Online First 10 September 2015



To cite: Korfei M, Skwarna S, Henneke I, *et al. Thorax* 2015;**70**: 1022–1032.

索

HDACi VPA.

Idiopathic pulmonary fibrosis (IPF) is a fatal lung disease characterised by destruction of alveolar

Key messages

What is the key question?

Although altered activity and expression of histone deacetylases (HDACs) have been reported in various pathological conditions, such as cancer or COPD, no data are currently available about the regional expression patterns of HDAC enzymes in idiopathic pulmonary fibrosis (IPF), a fatal lung disease for which no cure exists to date.

What is the bottom line?

This study shows for the first time that both myofibroblasts of fibroblast foci and bronchiolar basal cells of IPF lungs reveal 'cancer-like' overexpression of class I and class II HDAC enzymes, which may explain their progressive expansion as well as apoptosis resistance in this disease.

Why read on?

This study demonstrates for the first time that the pan-HDAC inhibitor LBH589/panobinostat, which is a proven antitumour agent, is capable of significantly downregulating collagen I biosynthesis and antiapoptotic genes in IPF fibroblast populations, and may thus potentially stop fibrotic remodelling in IPF.

architecture, abnormal fibroblast proliferation and excessive collagen deposition. Myofibroblasts are the primary collagen-producing cells in IPF, and their accumulation within pathological lesions called fibroblast foci (FF) is a key feature of IPF lungs.¹ Moreover, it has been suggested that IPF fibroblasts and myofibroblasts possess a cancer-like, 'malignant' phenotype,² with enhanced resistance to apoptosis due to increased expression of antiapoptotic and survival-related proteins, such as phosphatidylinositol-3-kinase- γ^3 or survivin.⁴ However, the precise mechanisms of such apoptosis resistance remain unclear.

BMJ

Structural changes of the chromatin, including alterations in the histone acetylation/deacetylation balance, have been suggested to underlie altered gene expression patterns in various pathological conditions, including cancer,⁵ cardiac hypertrophy⁶ and COPD.⁷ Histone deacetylases (HDACs) are enzymes that remove acetyl moieties from ɛ-N-acetylated lysine residues in histones, resulting in chromatin condensation and epigenetic repression of gene transcription. HDACs also catalyse deacetylation of many non-histone proteins, such as the tumour suppressor p53, resulting in inhibition of its (pro-apoptotic) transcriptional activity.⁸ ⁹ In line with their profound antiapoptotic activity. HDACs are upregulated in many types of cancer.^{5 8 9} In contrast, patients with COPD undergo a progressive reduction in total HDAC activity in lung tissue as a result of oxidative stress,⁷ which is associated with proteasomal degradation of many HDACs.¹⁰ HDACs are classified into four groups: class I (HDAC1, HDAC2, HDAC3 and HDAC8) are located primarily in the nuclei; class II (HDAC4, HDAC5, HDAC6, HDAC7, HDAC9 and HDAC10) are located in the cytoplasm but can shuttle to the nucleus; class III are NAD⁺-dependent enzymes and known as the sirtuins; class IV is represented by HDAC11 only.⁹ Class I, II and IV HDACs are Zn²⁺-dependent enzymes, and can thus be efficiently inhibited by chelating agents such as the hydroxamic acids vorinostat (SAHA) and panobinostat (LBH589).9 11 12 Importantly, these HDAC inhibitors (HDACi) have been reported as successful anticancer agents as they induce cell cycle arrest and apoptosis in cancer cells by increasing the acetylation status of chromatin and other nonhistone proteins.⁹ ¹¹ ¹²

Despite some reports about the involvement of altered HDAC activity in fibrotic lung fibroblasts and IPF,^{13–15} regulation and regional expression of HDACs have not been characterised in IPF. This study provides a comprehensive description of class I and class II HDAC expression patterns in the IPF lung. In addition, we can demonstrate that the profibrotic and 'malignant' phenotype of IPF fibroblasts can be greatly corrected by incubation with the pan-HDACi LBH589.

MATERIALS AND METHODS

Human lung tissue

Lung tissue samples were obtained from 26 patients with sporadic IPF (mean age \pm SD: 55.85 \pm 8.75 years; 5 females, 21 males), and 16 non-diseased control subjects (organ donors; mean age \pm SD: 48.92 \pm 15.84 years; 7 females, 8 males, 1 unknown). Explanted lungs were obtained from the Department of Thoracic Surgery, Vienna (W. Klepetko). All IPF diagnoses were made according to the American Thoracic Society/European Respiratory Society consensus criteria.¹ Four out of 26 patients with IPF were former smokers, 14 were never-smokers and from 8 patients the smoking status is unknown. With regard to control subjects, 3 out of 16 were former smokers, 7 were never-smokers and from 6 controls the smoking status is unknown.

Isolation of primary human lung fibroblasts

Primary human lung fibroblasts were isolated from explanted IPF (n=8) and control lungs (n=6) using an outgrowth technique as published.³ Experiments were carried out with IPF/ control fibroblasts between passages 3 and 4.

Cell culture experiments

At 95–99% confluency, IPF fibroblasts (n=6) were incubated for 30 h with the HDACi LBH589 (panobinostat, 85nM, Selleckchem) and valproic acid (VPA, 1.5 mM, Santa Cruz), and

as control experiment with the respective solvents (vehicle control) in the concentration 0.1% ethanol and 0.03% dimethyl sulfoxide. The dosages of HDACi were chosen according to published¹² ¹⁶ and own preliminary studies.

Methodology

Full details for quantitative immunoblotting, immunohistochemistry (IHC), semiquantitative RT-PCR and qRT-PCR (including primers) and statistics are available in the online supplementary material.

RESULTS

Upregulation of class I and class II HDACs in FF and abnormal bronchiolar epithelium in IPF lungs

Comparative immunoblot analysis of subpleural lung tissues from 26 patients with IPF versus 16 non-diseased control lungs revealed a significant upregulation of the class I HDACs HDAC1, HDAC2, HDAC3 and HDAC8 (see online supplementary figures S1A–D), as well as of the class IIa HDACs HDAC4, HDAC5, HDAC7 and HDAC9 (see online supplementary figures S2A–E), which comprises several isoforms (see online supplementary figures S2D, E). Protein expression of the class IIb HDAC10 was insignificantly upregulated (see online supplementary figure S2F). Though class IIb HDAC6 protein expression was not detectable by western blot, qRT-PCR indicated slightly increased *HDAC6* mRNA levels in subpleural IPF lung tissues compared with controls (see online supplementary figure S3).

Next, we employed IHC in order to identify the cellular distribution of these enzymes. A nuclear immunostaining for HDAC1 was observed in myofibroblasts of FF, which were identified by staining for alpha smooth muscle actin (a-SMA) (figure 1A, dashed arrows). Prominent nuclear expression of HDAC1 was also observed in overlying, hyperplastic bronchiolar basal cells (figure 1A, indicated by hashmark and cytoplasmic cytokeratin-5/KRT5 staining). Likewise, strong nuclear induction in FF and the overlying bronchiolar epithelium was encountered for the other class I members HDAC2/HDAC3 (see figure 1B and supplementary figures S4A/C). In addition, basal cell sheets of abnormal hyperplastic bronchioles in areas of bronchiolisation (indicated by KRT5 and nuclear p63 staining) showed strong nuclear expression of HDAC1 (figure 1C) and HDAC2/ HDAC3 (see online supplementary figure S5). HDAC2 and HDAC3 also revealed robust nuclear overexpression in ciliated bronchial cells in abnormal bronchiolar structures of IPF lungs (see online supplementary figures S4B and S6). Of note, neither HDAC1, HDAC2, nor HDAC3 showed a significant staining in type II alveolar epithelial cells (AECII) of IPF lungs (figure 1C, D, AECII indicated by arrows and proSP-C staining). In control lungs, HDAC1, HDAC2 and HDAC 3 expression was faint or nearly absent in AECII (figure 1E, indicated by arrows), and only weak expression of HDAC1, HDAC2 and HDAC3 could be encountered in basal and ciliated bronchial cells of normal bronchioles (see figure 1F and online supplementary figure S7).

Interestingly, robust cytoplasmic expression of HDAC3, but not of HDAC2, was detected in non-ciliated Clara cells of IPF and control lungs (see online supplementary figures S6 and S7B). The same observation was made in IHC for class I member HDAC8, which was expressed in general by bronchiolar epithelium in IPF and control lungs including basal, nonciliated and ciliated bronchial cells as well as by vascular smooth muscle cells (see online supplementary figures S8B, C). Moreover, cytoplasmic overexpression of HDAC8 was observed



Figure 1 Induction of class I histone deacetylases in fibroblast foci (FF) and abnormal bronchiolar epithelium of idiopathic pulmonary fibrosis (IPF) lungs. (A) Representative immunohistochemistry for cytokeratin-5 (KRT5), histone deacetylase 1 (HDAC1) and alpha-smooth muscle actin (α -SMA) in serial sections of IPF lung tissue. Strong nuclear staining for HDAC1 was predominantly observed in myofibroblasts of FF (indicated by dashed arrows and α -SMA-staining of a serial section) as well as in overlying hyperplastic bronchiolar basal cells (indicated by hashmark and KRT5 staining) of IPF lungs. (B) Representative immunohistochemistry for HDAC2, HDAC3 and KRT5 in serial sections of IPF lung tissue. In IPF, FF as well as overlying abnormal bronchiolar basal cells (indicated by hashmark and KRT5 expression) revealed strong nuclear induction of HDAC2 and HDAC3. (C) Representative immunohistochemistry for KRT5. HDAC1, prosurfactant protein C (proSP-C) and p63 in serial sections of IPF lung tissue. In IPF, bronchiolar basal cell sheets of hyperplastic bronchioles (positive for KRT5 and p63) indicated robust nuclear induction of HDAC1, whereas type II alveolar epithelial cells (AECII, indicated by arrows) indicated no expression of HDAC1. Only in very rare cases an expression of HDAC1 was observed in the IPF-AECII. AECII are identified by proSP-C immunostaining. (D) Representative immunohistochemistry for proSP-C, HDAC2, HDAC3 and KRT5 in serial sections of IPF lung tissue. IPF-AECII (indicated by arrows and proSP-C staining) did neither express HDAC2 nor HDAC3, which were found in the nucleus of bronchiolar basal cells in abnormal epithelia (positive for KRT5) as well as in some interstitial inflammatory cells (indicated by asterisk) in the very same IPF lung section. (E) Representative immunohistochemistry for proSP-C, HDAC1, HDAC2 and HDAC3 in serial sections of control lung tissue. No or only faint staining for HDAC1, HDAC2 and HDAC3 was observed in AECII of normal lungs; interstitial inflammatory cells (indicated by asterisk) of control lungs showed nuclear HDAC3 expression. (F) Representative immunohistochemistry for HDAC1 and KRT5 in serial sections of control lung tissue. Basal cells of normal bronchioles in control lungs (indicated by KRT5 staining) revealed no notable HDAC1 expression.

in myofibroblasts of FF (see online supplementary figure S8A), which is in line with previous reports describing that cytoplasmic HDAC8 associates with α -SMA and acts as an essential regulator of smooth muscle cell contractility.¹⁷

Concerning the cellular distribution of class II HDACs, HDAC4 was abundantly expressed by myofibroblasts of FF (figure 2A, dashed arrows) and in the overlying hyperplastic bronchiolar epithelium (marked by KRT5 staining, figure 2A), as well as in basal cell sheets of abnormal bronchiolar structures in IPF lungs (see online supplementary figure S9A), thereby indicating predominantly cytoplasmic localisation. A similar expression pattern was observed for class IIb HDAC10 (figure 2A, dashed arrows). Furthermore, HDAC4 was robustly expressed in the mature smooth muscle in areas of established fibrosis (indicated by α -SMA staining in figure 2B and online supplementary figure S10B), and co-localised with nuclear HDAC2 and survivin expression (figure 2B). Interestingly, ciliated bronchial cells of IPF lungs indicated strong nuclear expression of HDAC4 (indicated by arrowheads and FOXJ1 staining, figure 2C and supplementary figures S9B and S10C), which appeared to be weak in ciliated epithelium of normal bronchioles in control lungs (figure 2D). Notably, IPF-AECII did not indicate a significant expression of HDAC4 (figure 2C, arrows), whereas robust nuclear expression of HDAC4 was detected in the AECII of control lungs (figure 2D, arrows, and online supplementary figure S11A).

With regard to other class IIa HDAC members, HDAC7 immunostaining was strikingly pronounced in FF and bronchiolar epithelial cells of IPF lungs (figure 2E, F), but was faint in both IPF and donor AECIIs (see figure 2F, arrows, and online supplementary figure S11B). As expected, the expression of the other class IIa HDACs HDAC5 and HDAC9 isoform histone deacetylase-related protein (HDRP) was similarly distributed in IPF and control lungs (see online supplementary figures S12–S18).



Figure 2 Overexpression of class II histone deacetylases (HDACs) in fibroblast foci and abnormal bronchiolar epithelium of idiopathic pulmonary fibrosis (IPF) lungs. (A) Representative immunohistochemistry for p63, KRT5, HDAC4, alpha smooth muscle actin (α -SMA) and HDAC10 in serial sections of IPF lung tissue. In IPF, immunostaining for HDAC4 (cytoplasmic) and HDAC10 (nuclear and cytoplasmic) was observed in myofibroblasts of fibroblast foci (FF) (indicated by dashed arrows and α -SMA staining of serial sections) as well as in overlying hyperplastic bronchiolar basal cells (indicated by hashmarks, positive for p63 and KRT5). (B) Representative immunohistochemistry for KRT5, HDAC2, HDAC4, α -SMA and survivin in serial sections of IPF lung tissue. Established smooth muscle (indicated by α-SMA staining) in areas of dense 'older' fibrosis in IPF revealed cytoplasmic expression of HDAC4, with concomitant nuclear expression of HDAC2 and survivin. (C) Representative immunohistochemistry for KRT5, proSP-C, HDAC4, Forkhead box protein J1 (FOXJ1) and Clara cell protein 10 (CC10) in serial sections of IPF lung tissue. In IPF, strong nuclear staining for HDAC4 is observed in ciliated bronchial cells (indicated by arrowheads and FOXJ1 staining) and basal cells (positive for KRT5) of abnormal hyperplastic bronchioles. Type II alveolar epithelial cells (AECII, indicated by arrows and proSP-C staining) and Clara cells (positive for CC10) in IPF lungs did not express HDAC4. (D) Representative immunohistochemistry for proSP-C, HDAC4 and KRT5 in serial sections of control lung tissue. In the normal lung, HDAC4 was found to be expressed in the nucleus of AECII (indicated by arrows and proSP-C-staining), and basal cytoplasmic expression of HDAC4 was observed in bronchioles of control lungs. (E) Representative immunohistochemistry for p63, KRT5, HDAC7 and α -SMA in serial sections of IPF lung tissue. In IPF, the antibody for HDAC7 revealed strong cytoplasmic staining of myofibroblasts in FF as well as in overlying hyperplastic bronchiolar basal cells (indicated by hashmarks, positive for p63 and KRT5). (F) Representative immunohistochemistry for proSP-C and HDAC7 in serial sections of IPF lung tissue. AECII in areas of dense fibrotic tissue (indicated by arrows and proSP-C-staining) in IPF showed no or only faint immunostaining for HDAC7, whereas bronchial epithelium (BE) indicated robust cytoplasmic HDAC7 expression.

Next, cellular localisation of class IIb member HDAC6 was characterised in IPF and control lungs. HDAC6, which has been reported to function in aggresome-autophagy pathway,¹⁸ was the only HDAC with pronounced expression in AECII of IPF lungs. Moreover, HDAC6 cytoplasmic expression was

upregulated in IPF-AECII compared with control-AECIIs (see online supplementary figures \$19A,C, indicated by arrows). In addition, HDAC6 expression was increased in proSP-C-negative, AECII-like (or AECI-like) cells (see online supplementary figure \$20). Compared to HDAC7 expression, HDAC6 expression in myofibroblasts was mild (see online supplementary figure S19B). Notably, the ciliated epithelium in IPF and control lungs exhibited abundant cytoplasmic expression of HDAC6 (indicated by hashmarks in online supplementary figures S19A–C).

The IHC expression patterns for all described HDAC enzymes are summarised in table 1.

HDAC expression is upregulated in primary IPF fibroblasts

In accordance with previous reports showing persistence of a profibrotic phenotype of IPF-originating fibroblasts through several passages, we were able to detect a robust and significant upregulation of transcripts for the extracellular matrix (ECM)-associated genes COL1A1, COL3A1 and P4HTM in IPF (n=8) versus control (n=6) fibroblasts (see online supplementary figures S22A-D). ACTA2 appeared only slightly upregulated on mRNA level in IPF (see online supplementary figures S22A/ E). In line with mRNA expression, protein biosynthesis of procollagen I was significantly upregulated in IPF versus control fibroblasts (see online supplementary figure S22F). Moreover, primary IPF fibroblasts indicated a significant downregulation of CD90/Thy-1 antigen versus non-diseased controls (see online supplementary figure S22G), which is in line with many previous studies reporting loss of this cell surface glycoprotein in fibroblast populations of IPF lungs, especially in the myofibroblasts of FF.¹

Furthermore, RT-PCR indicated similar expression of *HDAC2* mRNA in IPF and control fibroblasts (figure 3A), but significant upregulation of mRNA transcripts for *HDAC3* (figure 3B), *HDAC4* (figure 3C), *HDAC5* (figure 3D), *HDAC7* (figure 3E) and *HDAC11* (figure 3F) in IPF fibroblasts. As anticipated by IHC analysis, quantitative immunoblotting revealed significant elevated protein expression of HDAC2 (figure 3G), HDAC3 (figure 3H), HDAC4 and HDAC7 (figure 3I), HDAC8 (figure 3J) and HDAC10 (figure 3K) in IPF versus control fibroblasts. HDAC9 protein expression was equal in IPF and control fibroblasts (see online supplementary figure S24). Additionally, IPF fibroblasts indicated reduced tubulin acetylation, which is indicative of increased expression of class IIb HDAC6 and HDAC10 (figure 3L).

Impact of HDACi treatment on survival and ECM production of IPF fibroblasts

Primary IPF fibroblasts (n=6) were treated with the pan-HDACi LBH589 (85nM) or the class I HDACi VPA (1.5 mM) or vehicle (Veh.) for 30 h, and analysed for cellular consequences. In LBH589 versus vehicle-treated IPF fibroblasts, mRNA expression of the matrix components *ACTA2* (figure 4A), *COL1A1* (figure 4B), *COL3A1* (figure 4C) and *FN* (figure 4D) were significantly decreased. Likewise, expression of the proliferation marker *CCND1* (figure 4E) and the

Cell type	HDAC expression pattern in IPF	HDAC expression pattern in control
Fibroblast foci	+++ HDAC1, HDAC2, HDAC3 (nucleus) +++ HDAC8, HDAC4, HDAC5, HDRP (cytoplasm) + HDAC6 (cytoplasm)* ++++ HDAC7 (cytoplasm) +++ HDAC10 (cytoplasm and nucleus)	-
Basal cell	+++ HDAC1, HDAC2, HDAC3 (nucleus) ++++ HDAC4 (cytoplasm and nucleus) +++ HDAC8, HDAC6 (cytoplasm) ++++ HDAC7 ++ HDAC5, HDRP (cytoplasm) +++ HDAC10 (cytoplasm and nucleus)	+ HDAC1 (nucleus) ++ HDAC4 (cytoplasm and nucleus) + HDAC7, HDRP (cytoplasm)
Ciliated bronchial cell	+++ HDAC2, HDAC3 (nucleus) +++ HDAC8, HDAC5 (cytoplasm) ++++ HDAC6*, HDAC7 (cytoplasm) ++++ HDAC4 (nucleus) ++ HDAC10 (cytoplasm)†	+ HDAC2, HDAC3 (nucleus) +++ HDAC8 (cytoplasm) + HDAC5, HDAC7, HDRP (cytoplasm) ++++ HDAC6 (cytoplasm)* ++ HDAC10 (cytoplasm)†
Clara cell	+++ HDAC3, HDAC8 (cytoplasm) ++ HDAC10 (cytoplasm)†	+++ HDAC3, HDAC8 (cytoplasm) ++ HDAC10 (cytoplasm)†
Unknown bronchial epithelial cell type	++++ HDRP (cytoplasm)‡	-
AECII	+ HDAC1 (nucleus) + HDAC5 (cytoplasm) +++ HDAC6 (cytoplasm)*	+ HDAC2, HDAC3 (nucleus) +++ HDAC4 (nucleus and cytoplasm) + HDAC5, HDAC7 (cytoplasm)
SMC	+++ HDAC2, HDAC3 (nucleus) +++ HDAC4 (cytoplasm) ++ HDAC7, HDAC8 (cytoplasm) + HDAC5 (cytoplasm)	+ HDAC4, HDAC7 (cytoplasm)
VSMC	++ HDAC1 (nucleus) +++ HDAC2, (nucleus) ++++ HDAC3 (nucleus+cytoplasm) +++ HDAC4, HDAC7, HDAC9, HDAC10 (cytoplasm) ++++ HDAC5 (cytoplasm) +++ HDAC8 (cytoplasm)	+ HDAC1, HDAC2, HDAC3 (nucleus) ++ HDAC4, HDAC5, HDAC7, HDAC9 (cytoplasm ++ HDAC8 (cytoplasm) + HDAC10 (cytoplasm and nucleus)

++++, very strong expression; +++, strong expression; +, very weak/faint expression; ++, weak-to-moderate expression; AECII, type II alveolar epithelial cell; HDAC, histone deacetylase; HDRP, histone deacetylase-related protein; IHC, immunohistochemistry; IPF, idiopathic pulmonary fibrosis; SMC, smooth muscle cell; VSMC, vascular smooth muscle cell. *See supplementary figures S19 and S20. tSee supplementary figure S21.

‡See supplementary-figures S16 and S17.

Interstitial lung disease



Figure 3 Expression analysis of histone deacetylases (HDACs) in primary fibroblasts isolated from idiopathic pulmonary fibrosis (IPF)–and non-diseased control lungs. (A–F) Reverse transcription-PCRs (RT-PCR) for *HDAC2* (A), *HDAC3* (B), *HDAC4* (C), *HDAC5* (D), *HDAC7* (E) and *HDAC11* (F) of human fibroblasts isolated from peripheral explanted lung tissue of patients with IPF (n=8) and non-diseased control lungs (control, n=6). Each PCR reaction was performed with 100 ng reverse-transcribed complementary DNA, followed by electrophoresis through a 2% (w/v) agarose gel containing ethidium bromide. Scanned agarose gels of indicated genes are shown in online supplementary figure S23. Band intensities of PCR products were densitometrically quantified, and mRNA expression of *HDAC* genes was normalised to the expression of glyceraldehyde 3-phosphate dehydrogenase (*GAPDH*). *p<0.05. (G–L) Representative immunoblots and quantitative immunoblot analysis of human fibroblasts isolated from peripheral lung tissue of patients with sporadic IPF (n=8) and non-diseased control lungs (control, n=6) for HDAC2 (G), HDAC3 (H), HDAC4 and HDAC7 (I), HDAC8 (J), HDAC10 (K), acetylated tubulin (L) and GAPDH or tubulin as loading controls. *p<0.05, **p<0.01, ***p<0.001. Densitometric ratios of the respective gene to *GAPDH*, as well as of the respective protein to GAPDH or tubulin, are depicted as a box-and-whisker diagram (*box* indicates 25th and 75th, *horizontal line* indicates the 50th percentile (median), and extensions above and below reflect extreme values); statistics were performed by Mann–Whitney test. x indicates this sample was not included for statistical analysis due to artefact.

survival gene *BIRC5* (figure 4F) was reduced, whereas mRNA levels for the proapoptotic genes *CIP1* (figure 4G) and *PUMA* (figure 4I) were increased. Notably, significant impairment of survival signalling was also observed in VPA-treated cells (figure 4E–G, I). However, a significant reduction of *P53* mRNA in response to LBH589 or VPA treatment (figure 4H) was observed, despite increases of p53 target genes *CIP1* and *PUMA* in both conditions. In line with apoptotic signalling, significant induction of the endoplasmic reticulum (ER)

stress-associated proapoptotic transcription factor *CHOP* was observed in LBH589 versus vehicle-treated cells, but not in VPA treatments (figure 4K). The mRNA expression of the ER stress marker ATF6 was slightly elevated in response to LBH589 treatment (figure 4J). Finally, VPA treatment leads to elevated transcription of the E2 conjugase *UBE2L6* (figure 4L) that has been described as part of the ubiquitination machinery for HDAC2, which is selectively degraded by the proteasome in response to VPA.¹⁶



Figure 4 Cellular signalling in primary idiopathic pulmonary fibrosis (IPF) fibroblasts in response to treatment with panobinostat or valproic acid (VPA). Primary IPF fibroblasts (n=6) were incubated for 30 h with vehicle (0.03% dimethyl sulfoxide, 0.1% ethanol), panobinostat (LBH589, 85 nmol) or VPA (1.5 mM). The effects of vehicle and histone deacetylase (HDACs) inhibitor treatments were analysed by semiquantitative reverse transcription-PCR (RT-PCR) for indicated genes. (A) *ACTA2*, (B) *COL1A1*, (C) *COL3A1*, (D) *FN*, (E) *CCND1*, (F) *BIRC5*, (G) *CIP1*, (H) *P53*, (I) *PUMA*, (J) *ATF6*, (K) *CHOP* and (L) *UBE2L6*. Each PCR reaction was performed with 100 ng reverse-transcribed complementary DNA, followed by electrophoresis through a 2% (w/v) agarose gel containing ethidium bromide. Scanned agarose gels of indicated genes are shown in online supplementary figure S25. Band intensities of PCR products were densitometrically quantified, and mRNA expression of indicated genes was normalised to the expression of glyceraldehyde 3-phosphate dehydrogenase (*GAPDH*). Data are presented as mean±SEM of the individual values of different treatments. *p<0.05, **p<0.01, ***p<0.001, LBH589 or VPA vs vehicle; *p<0.05, **p<0.01, ***p<0.001 VPA vs LBH589; by Dunn's multiple comparison test.

Next, gene expression of HDAC enzymes in response to HDACi treatment was analysed (see online supplementary figures S26A–N). Compared to vehicle-treated cells, LBH589 treatment resulted in reduced HDAC11 expression (see online supplementary figure S9L) and to marked suppression of HDAC7 (see online supplementary figure S9H), paralleled by a significant increase in HDAC3 (see online supplementary figure S9D) and SIRT2 transcripts (see online supplementary figureS9N).

On protein level (see figure 5 and online supplementary figure S27), IPF fibroblast lysates (n=4) indicated a significant increase of total acetylated histone H3 after treatment with LBH589 or VPA versus vehicle (figure 5A), whereas significant tubulin-acetylation was detected only in LBH589-treated cells (figure 5B) due to efficient inhibition of class IIb HDAC6 and HDAC10. In line with mRNA expression analysis, p21 protein was significantly increased (despite reduction of p53) in LBH589-treated IPF fibroblasts compared with vehicle control (figure 5C, D). Furthermore, western blot analyses displayed a decrease in proliferation marker phospho-histone H3 in LBH589-treated as well as VPA-treated IPF fibroblasts (figure 5E). Cyclin D1 was reduced only in LBH589-treated cells (figure 5F). Concomitantly, antiapoptotic proteins survivin (figure 5G) and Bcl-XL (figure 5H) were reduced, with parallel induction of the proapoptotic ER stress marker CHOP (figure 5I) and caspase-3 activation (figure 5J), in LBH589-treated, but not VPA-treated IPF fibroblasts. It has been already shown in cancer cells that ER stress plays a pivotal role in cell death mediated by LBH589, independent of p53.²⁰

In accordance with suppression of *HDAC7* gene expression in response to LBH589 treatment, protein expression of HDAC7 was almost absent under these conditions, but also reduced in VPA-treated IPF fibroblasts in comparison to vehicle control (see figure 5K and online supplementary figure S28; two different antibodies were tested for HDAC7 expression). In line with this interesting observation, the pan-HDACi SAHA has been shown to selectively suppress expression of *HDAC7* in several cell lines.²¹

Furthermore, VPA, but not LBH589 treatment, resulted in a significant downregulation of HDAC2 protein expression (figure 5L), presumably through E2-conjugase *UBE2L6*-mediated polyubiquitination and proteasomal degradation of HDAC2, a process that has been observed in several cell lines in response to VPA.¹⁶ In contrast, protein expression of HDAC3 was not altered by VPA, but was significantly upregulated in LBH589-treated versus vehicle-treated IPF fibroblasts (figure 5M).

Finally, all these changes were accompanied by significant reduction of collagen I biosynthesis in both LBH589-treated and VPA-treated IPF fibroblasts versus vehicle treatment, as assessed by immunoblotting for pro-forms (figure 5O) and mature forms of collagen I (figure 5P). Of note, VPA treatment did not reduce α -SMA expression in IPF fibroblasts (figure 5N).

DISCUSSION

The development, growth and aggressiveness of cancer cells have been associated with upregulated HDAC expression. Class I HDACs such as HDAC2 appear to be crucial for the regulation of proliferation and apoptosis resistance of cancer cells, in part



Figure 5 Analysis of acetylation status, apoptotic signalling and profibrotic protein expression in primary idiopathic pulmonary fibrosis (IPF) fibroblasts in response to treatment with panobinostat or valproic acid (VPA). Primary IPF fibroblasts (n=4) were incubated for 30 h with vehicle (0.03% dimethyl sulfoxide, 0.1% ethanol), panobinostat (LBH589, 85 nmol) or VPA (1.5 mM). Status of acetylation, apoptosis, histone deacetylase (HDAC) protein expression and profibrotic protein expression was analysed by quantitative immunoblotting. Representative immunoblots of indicated research targets are shown in online supplementary figure S27. In dependency of research target, histone H3, glyceraldehyde 3-phosphate dehydrogenase (GAPDH), β -actin or tubulin served as loading control. (A) Histone H3-acetyl K27, (B) acetylated tubulin, (C) p53, (D) p21, (E) phospho-histone H3, (F) cyclin-D1, (G) survivin, (H) Bcl-XI, (I) CHOP, (J) caspase-3, (K) HDAC7, (L) HDAC2, (M) HDAC3, (N) alpha-smooth muscle actin (α -SMA), (O) COL1A1 (pro-form) and (P) COL1A1 (mature form). Data are presented as mean±SEM of the individual values of different treatments. *p<0.05, **p<0.01, ***p<0.001, LBH589 or VPA vs vehicle; *p<0.05, **p<0.01, ***p<0.001 VPA vs LBH589; by Dunn's multiple comparison test.

through their ability to inhibit the tumour suppressor p53 and to promote expression of the proto-oncogene c-Myc.⁸ Several publications have also implicated many roles of class II HDACs in tumourigenesis: HDAC4 promotes growth of colon cancer cells via repression of p21.²² HDAC5 strongly interacts with the tumour suppressor RUNX3 and induces its degradation through deacetylation.²³ HDAC7 has been shown to protect from apoptosis by inhibiting c-Jun expression²⁴ and contributes to carcinogenesis by transcriptional activation of c-Myc.²⁵ However, little was known about expression and function of HDACs in IPF. In this paper, we show for the first time that nearly all class I and II HDAC enzymes are overexpressed and upregulated in IPF lung tissue. In detail, upregulation of HDACs was predominantly observed in myofibroblasts of FF as well as in abnormal bronchiolar basal cells in areas of bronchiolisation in IPF lungs, and co-localised with expression of survivin. These results are not surprising as key features in the pathogenesis of IPF include expansion of the population of fibroblasts and their differentiation into apoptosis-resistant myofibroblasts, which consecutively produce large amounts of ECM components.¹ While class I HDACs revealed strong nuclear induction in FF, class IIa HDACs HDAC4, HDAC5, HDAC7 and HDAC9 were predominantly expressed in the cytoplasm of myofibroblasts. The translocation of class IIa HDACs from the nucleus to the cytoplasm

associated with activation of transcription is factors myocyte-enhancer factor-2 and serum response factor, which activate myogenic genes such as ACTA2, and which are repressed by nuclear localisation of class IIa HDACs.¹⁴ Additional studies using RNAi technology identified HDAC4 as an HDAC that mediates transforming growth factor (TGF)-\u03b31-induced differentiation of normal lung fibroblasts into myofibroblasts. Also here, the cytoplasmic localisation of HDAC4 was required for activation of AKT signalling pathway, which is necessary for ACTA2 expression in response to TGF-β1.¹⁴ We suggest that 'cancer-like' upregulation of class I and II HDACs in IPF fibroblasts is associated with (i) increased fibroblast proliferation, (ii) fibroblast-to-myofibroblast differentiation and (iii) with the apoptosis-resistant phenotype of myofibroblasts. Another striking hallmark of IPF is the so-called 'bronchiolisation-process' that describes the aberrant progressive re-epithelialisation of damaged alveolar epithelium by proliferat-ing bronchiolar basal cells.^{1 26} Based on our data, we suggest that the exaggerated, proliferative character of basal cells in IPF may be in part due to overexpression of HDACs and that HDACs may govern the process of aberrant bronchiolisation in this disease. In line with this, luminal ciliated bronchial cells (which stem from the progenitor basal cells) indicated also robust overexpression of class I (especially HDAC2 and

A Class I+II HDACs are upregulated in fibroblasts and basal cells of IPF patient lungs



Figure 6 Aberrant histone deacetylase (HDAC) overexpression likely contributes to idiopathic pulmonary fibrosis (IPF) pathogenesis. (A) In the healthy lung with normal alveolar architecture, histone acetylation and deacetylation is balanced and contributes to appropriate gene transcription of pivotal cellular signalling pathways. The triggering event for IPF is alveolar epithelial cell (AEC) injury and death. AECII cells become hyperplastic and depleted of HDACs (except for HDAC6); endoplasmic reticulum stress and apoptosis are also present. In parallel, an aberrant upregulation of class I/II HDACs is observed in different fibroblast populations including myofibroblasts and bronchiolar basal cells. The contribution of injured IPF-AECIIs to the malignant HDAC overexpression in fibroblasts and basal cells in IPF remains to be elucidated. As a result of class I HDAC overexpression, we hypothesise that transcription of repressors of fibrosis is reduced in both fibroblast and basal cell populations of IPF lungs, thereby leading to aberrant activation of proliferative, profibrotic and apoptosis-resistant genes. Class II HDACs contribute to profibrotic gene expression by activating and maintaining cytoplasmic AKT phosphorylation. All together, this leads to abnormal fibroblast proliferation and myofibroblast generation. Alveolar spaces are infiltrated by myofibroblasts and activated resident fibroblasts, which begin to form fibroblast foci. Basal cells proliferate and migrate distally. As IPF progresses, alveolar spaces are progressively replaced by fibrotic tissue, smooth muscle and abnormal bronchiolar epithelium ('bronchiolisation'). (B) In vitro, pan-HDAC inhibition by LBH589/panobinostat efficiently inactivates primary profibrotic fibroblasts and myofibroblasts through downregulation of extracellular matrix (ECM) production on the one side and induction of apoptosis on the other side, and which thus may have a significant benefit for patients with IPF. (C) Legend.

HDAC3) and of all class II HDACs, whereas only faint or moderate expression was observed in the ciliated epithelium of nondiseased control lungs. Moreover, basal and luminal epithelial cells near areas of active fibrotic remodelling and aberrant bronchiolar proliferation or of abnormal IPF bronchioles that often appear hyperplastic and larger in size are suspected to contribute to the abnormal, exaggerated production of profibrotic signals in IPF²⁶ and that thus may require enhanced overexpression of 'cancer-like' HDACs. Of note, expression of many HDAC enzymes seemed to be virtually absent or was only faint in normal bronchiolar structures as well as AECII (or rather in the whole lung tissue) of non-diseased control lungs, presumably due to their crucial intrinsic activity in cell proliferation and apoptosis inhibition,^{5 8 9} which justifies an only low and basal expression in normal, healthy cells.

The class IIa HDACs HDAC4 and HDAC7 were the only HDACs with moderate to robust expression in ciliated epithelium and AECII of non-diseased control lungs. However, we can only speculate about the function of both these HDACs in the normal AECII. HDAC4 and HDAC7 have been both reported to regulate the functional activity of HIF-1α in various cell types in vitro because loss of each of them resulted in reduced HIF-1a transcriptional activity of adaptive responses to hypoxia.^{27 28} For alveolar-epithelial expressed HIF-1a, a surprising role in lung protection during acute lung injury (ALI) has recently been reported in which normoxic HIF-1a stabilisation and HIF-1\alpha-dependent control of alveolar-epithelial glucose metabolism attenuated lung inflammation during ALI induced by stretch conditions in vitro or by ventilator-induced lung injury in vivo.²⁹ Additionally, HIF-1 α has been reported to be essential for proper lung maturation and surfactant production in the newborn lung of mice.³⁰ Based on these studies, we can presume that HDAC4 and HDAC7 might play a crucial role in maintaining AECII homeostasis through regulation of HIF-1α.

Though, and in marked contrast to bronchiolar epithelial cells in IPF, AECII of IPF lungs did not reveal significant expression of class I and II HDACs (which were in part significantly expressed in AECII of normal lungs, such as HDAC4 and HDAC7 as described above), possibly due to pro-apoptotic ER and oxidative stress in IPF-AECII,^{31–33} which in turn lead to proteasomal degradation of many HDACs. Moreover, HDAC4 is a death substrate, cleaved by caspase-3 during apoptotic cell death,³⁴ and HDAC7 is the best caspase-8 substrate described to date.³⁵

Importantly, the class IIb member HDAC6 was the only HDAC expressed in IPF-AECII, but not in AECII of normal lungs. As a tubulin deacetylase, HDAC6 is involved in aggresome formation and autophagic clearance of protein aggregates in response to increased misfolding,¹⁸ and its induction in AECII can be due to severe alveolar epithelial ER stress in IPF.³¹⁻³³ Additionally, HDAC6 deacetylates antioxidants PRDX1 and PRDX2, which impairs their function to reduce H_2O_2 , thereby promoting apoptosis.³⁶ Despite its reported pro-apoptotic and pathological function, it seemed to be an intrinsic normality of ciliated bronchial cells of IPF as well as non-diseased control lungs to robustly express HDAC6 in the cytoplasm, as shown by our IHC studies. Of note, autophagy is also critical for cell survival because unwanted intracellular materials are degraded, and HDAC6 controls autophagosome maturation through its deacetylase activity.³⁷ Indeed, it was shown that cells deficient in HDAC6 failed to form autophagosomes and to clear misfolded protein aggregates, and were hypersensitive to the accumulation of malfolded proteins.³⁷ On the other hand, inhibition of HDAC6 deacetylase activity by pan-HDACi such as SAHA or trichostatin-A has been shown to ameliorate disease progression and neurodegeneration in rodent models of Huntington's disease and amyotrophic lateral sclerosis.³⁸ ³⁹ Anyway, the exact role of HDAC6 in the ciliated lung epithelium and the impact of its induction in IPF-AECII remain still elusive and have to be further investigated.

More than 10 HDACi have already been examined in clinical trials as cancer drugs and only minor side effects have been observed.⁹ HDACi possess the ability to selectively induce apoptosis in tumour cells, whereas normal cells are relatively resistant to HDACi-induced cell death.⁹ ¹¹ ⁴⁰ In addition to antitumour activities, HDACi have been shown to exert beneficial effects on several diseases such as cardiac hypertrophy,⁶ inflammatory bowel disease⁴¹ and above-mentioned neurodegenerative diseases.³⁸ ³⁹ The present study suggests that HDACi such as LBH589 may also be useful for the treatment of IPF. Interfering

Interstitial lung disease

with the generation of myofibroblast foci to decrease production of ECM components is important in treating IPF because the increase in the number of these fibrotic foci has been associated with disease progression and a worsened prognosis.⁴² We demonstrated that LBH589 downregulated significantly the mRNA expression of ACTA2 and ECM genes COL1A1, COL3A1 and FN in primary IPF fibroblasts and thus interfered with fibroblast-to-myofibroblast differentiation. In addition, LBH589 efficiently reduced the protein biosynthesis of the large collagen α -1 chains (COL1A1) in IPF fibroblasts. The reduction in ECM synthesis was accompanied by growth inhibition as shown by immunoblotting for p-histone H3 and cyclin-D1. In parallel, LBH589 suppressed expression of the survival-related HDAC7 gene and downregulated expression of antiapoptotic genes BCL2L1/Bcl-XL and BIRC5/survivin in IPF fibroblasts, and led to induction of ER stress and apoptosis, thereby resulting in efficient inactivation of ECM producing and 'malignant appearing' IPF fibroblasts. Moreover, in renal fibrosis, it was demonstrated in a murine model of unilateral ureteral obstruction that trichostatin-A led simultaneously to inactivation of renal interstitial fibroblasts and inhibition of renal tubular epithelial cell death,⁴³ thus adding further experimental proof to use pan-HDACi for the treatment of IPF. Furthermore, it should be noted that the treatment of IPF fibroblasts with VPA, which has been described as a weak class I HDACi, led also to a significant reduction in collagen I biosynthesis. Additionally, VPA treatment induced significant downregulation of HDAC2 protein levels in IPF fibroblasts, presumably through UBE2L6-mediated polyubiquitination and proteasomal degradation of HDAC2.¹⁶ Because overexpression of HDAC2 is a classical 'cancer condition', drugs that interfere with HDAC2 protein turnover in myofibroblasts are of great importance.

The exact targets of HDACs in the setting of fibrosis are yet not clear. Similar to cancer cells, in which 'aberrant transcriptional repression' has been described as a result of class I HDAC overexpression,^{5 8 11 16} we suggest that target genes of class I HDACs in lung fibroblasts might be repressors of antiapoptotic and profibrotic gene expression (which are silenced in the setting of fibrosis due to class I HDAC upregulation). The contribution of class IIa HDACs to lung fibrosis is presumably antiapoptosis^{22–25} as well as maintaining of cytoplasmic AKT phosphorylation/activation and consequent promotion of profibrotic gene expression.¹⁴

In conclusion, based on the results of our study, HDACs are novel molecular targets for IPF therapy, and pan-HDACi such as LBH589 are promising therapeutic agents for the treatment of IPF, even though their exact targets and mechanisms of action in the setting of IPF must be further investigated. The main observations and suggestions from this study are summarised in figure 6.

Author affiliations

¹Department of Internal Medicine, Justus-Liebig-University Giessen, Giessen, Germany

²Universities of Giessen and Marburg Lung Center (UGMLC), Member of the German Center for Lung Research (DZL)

³Department of Medicine, Section of Pulmonary Diseases, Critical Care and Environmental Medicine, Tulane University Health Science Center, New Orleans, Louisiana, USA

⁴Excellence Cluster Cardio-Pulmonary System (ECCPS), Giessen, Germany

⁵Department of Thoracic Surgery, Vienna General Hospital, Vienna, Austria

⁶CHU Paris Nord-Val de Seine, Hôpital Xavier Bichat-Claude Bernard, Paris, France ⁷Department of Lung Development and Remodeling, Max-Planck-Institute for Heart and Lung Research, Bad Nauheim, Germany

⁸Institute of Pathology and Cytology, Wetzlar, Germany

⁹Department of Toxicology, University Medical Center, Mainz, Germany

¹⁰Agaplesion Lung Clinic Waldhof Elgershausen, Greifenstein, Germany

¹¹European IPF Network and European IPF Registry

Interstitial lung disease

Acknowledgements We thank Gabriele Dahlem and Susanna Ziegler for isolation of primary fibroblasts from explanted IPF and non-diseased control lungs.

Contributors MK, OHK and AG designed experiments and wrote the manuscript. MK, SS, IH, BAMK, OK, ShS, LF, CR, DvdB, PM, and WK carried out the experimental work. BAMK, SB, BC, SSP and WS served as key advisors and critically read the manuscript. OHK and AG equally contributed to this manuscript.

Competing interests None declared.

Patient consent Obtained.

Ethics approval The study protocol was approved by the Ethics Committee of the Justus-Liebig-University School of Medicine (no. 31/93, 29/01, and no. 111/08: European IPF Registry).

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- American Thoracic Society. Idiopathic pulmonary fibrosis: diagnosis and treatment. International consensus statement. American Thoracic Society (ATS), and the European Respiratory Society (ERS). Am J Respir Crit Care Med 2000;161:646–64.
- Vancheri C. Common pathways in idiopathic pulmonary fibrosis and cancer. Eur Respir Rev 2013;22:265–72.
- 3 Conte E, Gili E, Fruciano M, et al. PI3K p110gamma overexpression in idiopathic pulmonary fibrosis lung tissue and fibroblast cells: in vitro effects of its inhibition. *Lab Invest* 2013;93:566–76.
- 4 Sisson TH, Maher TM, Ajayi IO, et al. Increased survivin expression contributes to apoptosis-resistance in IPF-fibroblasts. Adv Biosci Biotechnol 2012;3:657–64.
- 5 Lafon-Hughes L, Di Tomaso MV, Mendez-Acuna L, et al. Chromatin-remodelling mechanisms in cancer. Mutat Res 2008;658:191–214.
- 6 Backs J, Olson EN. Control of cardiac growth by histone acetylation/deacetylation. *Circ Res* 2006;98:15–24.
- 7 Ito K, Ito M, Elliott WM, et al. Decreased histone deacetylase activity in chronic obstructive pulmonary disease. New Engl J Med 2005;352:1967–76.
- 8 Brandl A, Wagner T, Uhlig KM, et al. Dynamically regulated sumoylation of HDAC2 controls p53 deacetylation and restricts apoptosis following genotoxic stress. J Mol Cell Biol 2012;4:284–93.
- 9 Xu WS, Parmigiani RB, Marks PA. Histone deacetylase inhibitors: Molecular mechanisms of action. *Oncogene* 2007;26:5541–52.
- 10 Sundar IK, Yao H, Rahman I. Oxidative stress and chromatin remodelling in chronic obstructive pulmonary disease and smoking-related diseases. *Antioxid Redox Signal* 2013;18:1956–71.
- 11 Richon VM, Garcia-Vargas J, Hardwick JS. Development of vorinostat: current applications and future perspectives for cancer therapy. *Cancer Lett* 2009;280:201–10.
- 12 Anne M, Sammartino D, Barginear MF, et al. Profile of panobinostat and its potential for treatment in solid tumors: an update. Onco Targets Ther 2013;6:1613–24.
- 13 Coward WR, Watts K, Feghali-Bostwick CA, et al. Defective histone acetylation is responsible for the diminished expression of cyclooxygenase 2 in idiopathic pulmonary fibrosis. *Mol Cell Biol* 2009;29:4325–39.
- 14 Guo W, Shan B, Klingsberg RC, et al. Abrogation of TGF-beta1-induced fibroblast-myofibroblast differentiation by histone deacetylase inhibition. Am J Physiol Lung Cell Mol Physiol 2009;297:L864–870.
- 15 Sanders YY, Hagood JS, Liu H, et al. Histone deacetylase inhibition promotes fibroblast apoptosis and ameliorates pulmonary fibrosis in mice. Eur Respir J 2014;43:1448–58.
- 16 Krämer OH, Zhu P, Ostendorff HP, et al. The histone deacetylase inhibitor valproic acid selectively induces proteasomal degradation of HDAC2. EMBO J 2003;22:3411–20.
- 17 Waltregny D, Glenisson W, Tran SL, *et al*. Histone deacetylase HDAC8 associates with smooth muscle alpha-actin and is essential for smooth muscle cell contractility. *FASEB J* 2005;19:966–8.
- 18 Lam HC, Cloonan SM, Bhashyam AR, et al. Histone deacetylase 6-mediated selective autophagy regulates COPD-associated cilia dysfunction. J Clin Invest 2013;123:5212–30.

- 19 Hagood JS, Prabhakaran P, Kumbla P, *et al.* Loss of fibroblast Thy-1 expression correlates with lung fibrogenesis. *Am J Pathol* 2005;167:365–79.
- 20 Montalbano R, Waldegger P, Quint K, et al. Endoplasmic reticulum stress plays a pivotal role in cell death mediated by the pan-deacetylase inhibitor panobinostat in human hepatocellular cancer cells. *Transl Cncol* 2013;6:143–57.
- 21 Dokmanovic M, Perez G, Xu W, et al. Histone deacetylase inhibitors selectively suppress expression of HDAC7. *Mol Cancer Ther* 2007;6:2525–34.
- 22 Wilson AJ, Byun DS, Nasser S, et al. HDAC4 promotes growth of colon cancer cells via repression of p21. Mol Biol Cell 2008;19:4062–75.
- 23 Jin YH, Jeon EJ, Li QL, et al. Transforming growth factor-beta stimulates p300-dependent RUNX3 acetylation, which inhibits ubiquitination-mediated degradation. J Biol Chem 2004;279:29409–17.
- 24 Ma C, D'Mello SR. Neuroprotection by histone deacetylase-7 (HDAC7) occurs by inhibition of c-jun expression through a deacetylase-independent mechanism. J Biol Chem 2011;286:4819–28.
- 25 Zhu C, Chen Q, Xie Z, *et al.* The role of histone deacetylase 7 (HDAC7) in cancer cell proliferation: Regulation on c-myc. *J Mol Med* 2011;89:279–89.
- 26 Chilosi M, Poletti V, Murer B, et al. Abnormal re-epithelialization and lung remodelling in idiopathic pulmonary fibrosis: the role of deltaN-p63. Lab Invest 2002;82:1335–45.
- 27 Wang Z, Qin G, Zhao TC. HDAC4: mechanism of regulation and biological functions. *Epigenomics* 2014;6:139–50.
- 28 Witt O, Deubzer HE, Milde T, et al. HDAC family: what are the cancer relevant targets? Cancer Lett 2009;277:8–21.
- 29 Eckle T, Brodsky K, Bonney M, et al. HIF1A reduces acute lung injury by optimizing carbohydrate metabolism in the alveolar epithelium. PLoS Biol 2013;11:e1001665.
- 30 Saini Y, Harkema JR, LaPres JJ. HIF1alpha is essential for normal intrauterine differentiation of alveolar epithelium and surfactant production in the newborn lung of mice. J Biol Chem 2008;283:33650–7.
- 31 Lawson WE, Crossno PF, Polosukhin VV, et al. Endoplasmic reticulum stress in alveolar epithelial cells is prominent in IPF: association with altered surfactant protein processing and herpesvirus infection. Am J Physiol Lung Cell Mol Physiol 2008;294:L1119–26.
- 32 Korfei M, Ruppert C, Mahavadi P, et al. Epithelial endoplasmic reticulum stress and apoptosis in sporadic idiopathic pulmonary fibrosis. Am J Respir Crit Care Med 2008;178:838–46.
- 33 Cha SI, Ryerson CJ, Lee JS, et al. Cleaved cytokeratin-18 is a mechanistically informative biomarker in idiopathic pulmonary fibrosis. *Respir Res* 2012;13:105.
- 34 Paroni G, Mizzau M, Henderson C, et al. Caspase-dependent regulation of histone deacetylase 4 nuclear-cytoplasmic shuttling promotes apoptosis. *Mol Biol Cell* 2004;15:2804–18.
- 35 Scott FL, Fuchs GJ, Boyd SE, et al. Caspase-8 cleaves histone deacetylase 7 and abolishes its transcription repressor function. J Biol Chem 2008;283:19499–510.
- 36 Parmigiani RB, Xu WS, Venta-Perez G, et al. HDAC6 is a specific deacetylase of peroxiredoxins and is involved in redox regulation. Proc Natl Acad Sci USA 2008;105:9633–8.
- 37 Lee JY, Koga H, Kawaguchi Y, et al. HDAC6 controls autophagosome maturation essential for ubiquitin-selective quality-control autophagy. *EMBO Journal* 2010;29:969–80.
- 38 Hockly E, Richon VM, Woodman B, et al. Suberoylanilide hydroxamic acid, a histone deacetylase inhibitor, ameliorates motor deficits in a mouse model of Huntington's disease. Proc Natl Acad Sci USA 2003;100:2041–6.
- 39 Petri S, Kiaei M, Kipiani K, et al. Additive neuroprotective effects of a histone deacetylase inhibitor and a catalytic antioxidant in a transgenic mouse model of amyotrophic lateral sclerosis. *Neurobiol Dis* 2006;22:40–9.
- 40 Qiu L, Burgess A, Fairlie DP, et al. Histone deacetylase inhibitors trigger a G2 checkpoint in normal cells that is defective in tumor cells. *Mol Biol Cell* 2000;11:2069–83.
- 41 Tao R, de Zoeten EF, Ozkaynak E, *et al*. Deacetylase inhibition promotes the generation and function of regulatory T cells. *Nat Med* 2007;13:1299–307.
- 42 King TE Jr, Schwarz MI, Brown K, et al. Idiopathic pulmonary fibrosis: relationship between histopathologic features and mortality. Am J Respir Crit Care Med 2001;164:1025–32.
- 43 Pang M, Kothapally J, Mao H, et al. Inhibition of histone deacetylase activity attenuates renal fibroblast activation and interstitial fibrosis in obstructive nephropathy. Am J Physiol Renal Physiol 2009;297:F996–F1005.