

Efficacy and safety outcomes of recanalisation procedures in patients with acute symptomatic pulmonary embolism: systematic review and network meta-analysis

David Jimenez,^{1,2} Carlos Martin-Saborido,³ Alfonso Muriel,⁴ Javier Zamora,⁴ Raquel Morillo,^{1,2} Deisy Barrios,^{1,2} Frederikus A Klok,⁵ Menno V Huisman,⁵ Victor Tapson,⁶ Roger D Yusen⁷

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/thoraxjnl-2017-210040>).

For numbered affiliations see end of article.

Correspondence to

Dr David Jimenez, Respiratory Department, Ramón y Cajal Hospital, Madrid 28034, Spain; djimenez.hrc@gmail.com

DJ and CM-S contributed equally.

Received 23 January 2017
Revised 13 October 2017
Accepted 23 October 2017
Published Online First
13 November 2017

ABSTRACT

Background We aimed to review the efficacy and safety of recanalisation procedures for the treatment of PE.

Methods We searched PubMed, the Cochrane Library, EMBASE, EBSCO, Web of Science and CINAHL databases from inception through 31 July 2015 and included randomised clinical trials that compared the effect of a recanalisation procedure versus each other or anticoagulant therapy in patients diagnosed with PE. We used network meta-analysis and multivariate random-effects meta-regression to estimate pooled differences between each intervention and meta-regression to assess the association between trial characteristics and the reported effects of recanalisation procedures versus anticoagulation.

Results For all-cause mortality, there were no significant differences in event rates between any of the recanalisation procedures and anticoagulant treatment (full-dose thrombolysis: OR 0.60; 95% CI 0.36 to 1.01; low-dose thrombolysis: 0.47; 95% CI 0.14 to 1.59; and catheter-associated thrombolysis: 0.31; 95% CI 0.01 to 7.96). Full-dose thrombolysis increased the risk of major bleeding (2.00; 95% CI 1.06 to 3.78) compared with anticoagulation. Catheter-directed thrombolysis was associated with the lowest probability of dying (surface under the cumulative ranking curve (SUCRA), 0.67), followed by low-dose thrombolysis (SUCRA, 0.66) and full-dose thrombolysis (SUCRA, 0.55). Similarly, low-dose thrombolysis was associated with the lowest probability of major bleeding (SUCRA, 0.61), followed by catheter-directed thrombolysis (SUCRA, 0.54) and full-dose thrombolysis (SUCRA, 0.17). The results were similar in sensitivity analyses based on restricting only to studies in haemodynamically stable patients with PE.

Conclusions In the treatment of PE, recanalisation procedures do not seem to offer a clear advantage compared with standard anticoagulation. Low-dose thrombolysis was associated with the lowest probability of dying and bleeding.

Trial registration number PROSPERO CRD42015024670.

INTRODUCTION

Although most patients with acute PE have an uncomplicated clinical course while undergoing

Key messages

What is the key question?

► For treatment of acute PE, the benefits and risks of the different recanalisation procedures (ie, full-dose systemic thrombolysis, reduced-dose systemic thrombolysis or catheter-directed thrombolysis) versus each other lack clarity.

What is the bottom line?

► Compared with standard anticoagulation, recanalisation procedures had a similar risk of all-cause mortality, and full-dose thrombolysis was associated with an increased risk of major bleeding.

Why read on?

► Low-dose thrombolysis was associated with the lowest probability of dying and bleeding.

standard anticoagulation treatment, the overall short-term mortality rate is still significant.^{1,2} Death from acute PE usually occurs before or soon after hospital admission.^{3,4}

There have been two main treatments for acute PE, anticoagulant therapy alone or systemic thrombolytic therapy.⁵ Most patients presenting to the hospital with PE have normal blood pressure, normal right ventricular function and a low clinical severity score and therefore have a very low short-term mortality with prompt initiation of anticoagulation. Although systemic thrombolysis has angiographic and haemodynamic benefits for patients with acute PE, compared with standard therapy, it markedly increases major bleeding, including intracranial and fatal bleeding.⁶ Consequently, systemic thrombolytic therapy is usually reserved for patients with PE with haemodynamic instability.⁷ The ability to actively remove emboli in patients with acute PE without increasing bleeding would be an important advance. Low-dose systemic thrombolysis and catheter-based thrombolytic therapy require only a fraction of the systemic fibrinolytic dose, and this dose reduction might improve the safety of thrombolysis for PE. A common problem in evaluating the efficacy of these interventions is the lack of trials (or



► <http://dx.doi.org/10.1136/thoraxjnl-2017-211141>



To cite: Jimenez D, Martin-Saborido C, Muriel A, *et al.* *Thorax* 2018;**73**:464–471.



a paucity of available trials) that directly compare these interventions. As a result, no meta-analysis has comprehensively compared the effect of a recanalisation procedure versus each other in patients diagnosed with acute symptomatic PE.

The primary aim of our study was to perform a network meta-analysis of randomised controlled trials (RCTs) for treatment of acute PE to obtain a better estimate of the benefits and risks of the different recanalisation procedures (ie, full-dose systemic thrombolysis, reduced-dose systemic thrombolysis or catheter-directed thrombolysis) versus each other or anticoagulant therapy.

METHODS

Data sources and searches

This systematic review is reported according to the Preferred Reporting Items for Systematic Reviews and Meta-analyses statement extension for network meta-analysis and was conducted following an a priori established protocol registered with PROSPERO.⁸ We searched PubMed, the Cochrane Library, EMBASE, EBSCO, Web of Science and CINAHL databases. Each database was searched from its inception date to 31 July 2015. Conference abstracts were included in our search. The retrieved articles were examined to eliminate potential duplicates or overlapping data. No limits or language restriction were applied during the search. The RCTs were identified using the Cochrane Collaboration highly sensitive search strategy (sensitivity-maximising and precision-maximising version).⁹ The search string was: (1) pulmonary embolis*; (2) thrombolysis OR thrombolytic therapy OR streptokinase OR urokinase OR tenecteplase OR alteplase OR desmoteplase OR tissue plasminogen activator OR clot-dissolving medication; (3) search strings 1 AND 2. We also hand searched the references of relevant articles for additional clinical trials not identified by the electronic search and contacted experts. Finally, we searched ClinicalTrials.gov for information on clinical trials that were terminated but unpublished. The planned analysis was registered at the PROSPERO international prospective register of systematic reviews on 20 July 2015 (CRD42015024670).

Study selection

One reviewer (DJ) performed the database search and initial screening of titles and abstracts. Two investigators (DJ and RM) independently carried out full-text screening of all eligible articles. We included a study if participants were patients with acute symptomatic PE objectively diagnosed with standard imaging techniques and received anticoagulant therapy; the intervention was treatment with a recanalisation procedure (ie, full-dose systemic thrombolysis, reduced-dose systemic thrombolysis or catheter-directed thrombolysis); the comparison group was either treatment with a different recanalisation procedure or no recanalisation treatment (ie, the patients received standard anticoagulation); it was an RCT; and it reported mortality outcomes. Observational studies and trials without a control group were excluded.

Data extraction and quality assessment

Two reviewers (DJ and RM) independently extracted data onto a computer spreadsheet, with discrepancies resolved by consensus. Extracted data included first author, year of publication, type of intervention and control group, number of patients, patient characteristics and duration of follow-up. The primary outcomes were all-cause mortality and major bleeding, as defined by the study protocol. Secondary outcomes were risk of intracranial

haemorrhage (ICH) and recurrent embolism. The occurrence of these outcomes was abstracted according to the intention-to-treat population for individual trials. The outcomes data from the first available time point identified as a primary end point from each trial were incorporated into our primary analysis. Each study was graded for potential bias into low, high and unclear according to the Cochrane Collaboration handbook.¹⁰

Data synthesis and analysis

Separate meta-analyses of direct evidence only (pairwise meta-analyses) were performed using DerSimonian and Laird random-effects model to estimate pooled ORs and 95% CIs.¹¹ Forest plots were created for each outcome. When there were no events in one treatment group, we used a 0.5 continuity correction. Heterogeneity was assessed using the estimated between-study variance (τ^2), Cochran χ^2 test and the I^2 statistic.¹²

Because there are few trials making head-to-head comparisons between recanalisation procedures, we performed a network meta-analysis. Unlike traditional meta-analyses, this method has the advantage of allowing trials comparing recanalisation procedures with some other common treatment (eg, placebo) to be incorporated into the analysis, thus increasing power and enabling a better comparison of recanalisation therapies to be made.¹³ We used multivariate, random-effects meta-regressions to perform each analysis using the network family of commands in Stata.¹⁴ We evaluated inconsistency between direct and indirect sources of evidence by comparison of the fit and parsimony of consistency and inconsistency models and by calculation of the difference between direct and indirect estimates of a specific treatment effect ('loop-specific approach'). The relative ranking of recanalisation interventions on primary and secondary outcomes was presented as their surface under the cumulative ranking curve (SUCRA) probabilities, which represent their likelihood of being ranked best.¹⁵ In this study, higher SUCRA scores reflect lower associated all-cause mortality and bleeding events. We estimated the probability of each treatment being the best by averaging 10 000 Monte Carlo replications. The level of statistical significance was set at $P < 0.05$ and all statistical tests were 2-sided.

We performed some sensitivity analyses to assess the robustness of the findings. These were based on (1) restricting only to studies in patients with haemodynamically stable PE; (2) restricting only to trials where the mean age of participants in the thrombolytic group was > 65 years; and (3) alternative statistical model (frequentist approach using a random-effects inconsistency model).

RESULTS

From a total of 930 unique studies identified using the search strategy, 22 RCTs (2494 patients) were included in the network meta-analysis (online supplementary efigure 1). These included 16 trials comparing full-dose thrombolysis to no thrombolysis (2016 patients),^{6 16–30} 1 comparing low-dose thrombolysis to no thrombolysis (121 patients),³¹ 1 comparing ultrasound-assisted catheter-directed thrombolysis with no thrombolysis (59 patients)³² and 4 comparing full-dose thrombolysis with low-dose thrombolysis (298 patients).^{33–36} The available direct comparisons and network of trials are shown in figure 1 and online supplementary efigures 2–4.

Characteristics of included studies

The RCTs included in the network meta-analysis are summarised in table 1. Overall, these 22 trials were reported between 1970




Figure 1 Network of included studies with available direct comparisons for all-cause mortality.

and 2014 and included 2494 participants. The mean study sample size was 113 participants, ranging from 8 to 1005 patients. The baseline characteristics of patients included in these trials are described in [table 1](#). The primary outcome (all-cause mortality) was reported in all studies.

Direct meta-analysis

Results of direct pairwise meta-analysis are summarised in [table 2](#) and online supplementary efigures 5–8. All interventions were associated with a non-significant reduction of all-cause mortality (full-dose thrombolysis: OR 0.64; 95% CI 0.37 to 1.09; low-dose thrombolysis: 0.32, 0.03 to 3.13; catheter-directed thrombolysis: 0.31, 0.01 to 7.96); full-dose thrombolysis was not superior to low-dose thrombolysis (1.04, 0.24 to 4.41). Full-dose thrombolytic therapy was significantly associated with a greater risk of major bleeding (2.39, 1.44 to 3.95) and ICH (3.66, 1.13 to 11.86) compared with anticoagulant therapy (online supplementary efigures 6 and 7), whereas low-dose thrombolysis showed a non-significant benefit in terms of major bleeding and ICH compared with full-dose thrombolysis ([table 2](#)). All outcomes were associated with negligible heterogeneity ($I^2 < 12\%$).

Network meta-analysis: primary outcomes

In network meta-analysis, compared with anticoagulation alone, full-dose thrombolysis was associated with an OR of 0.60 (95% CI 0.36 to 1.01), low-dose thrombolysis with an OR of 0.47 (95% CI 0.14 to 1.59) and catheter-directed thrombolysis with an OR of 0.31 (95% CI 0.01 to 7.96) for dying ([figure 2](#)). When recanalisation treatments were compared, none of comparisons reached conventional level of statistical significance ([figure 2](#)). In network meta-analysis, compared with anticoagulation alone, full-dose thrombolysis was associated with an OR of 2.00 (95% CI 1.06 to 3.78), low-dose thrombolysis with an OR of 0.90 (95% CI 0.25 to 3.21) and catheter-directed thrombolysis with an OR of 0.97 (95% CI 0.02 to 56.03) for bleeding ([figure 2](#)). Again, when recanalisation treatments were compared for bleeding, none of comparisons reached conventional level of statistical significance ([figure 2](#)).

Network meta-analysis suggested that catheter-directed thrombolysis was associated with the lowest probability of dying (SUCRA, 0.67), followed by low-dose thrombolysis (SUCRA, 0.66) and full-dose thrombolysis (SUCRA, 0.55) ([figure 3](#)). Similarly, low-dose thrombolysis was associated with the lowest

Table 1 Characteristics of included randomised clinical trials

Source	Patients (n)	Intervention*	Control*	High-risk PE included	Age, mean (range or SD), years	Follow-up, d	Male, n (%)	All-cause mortality	Major bleeding	ICH	Recurrent VTE
Meyer <i>et al</i> ⁶	1005	Tenecteplase (30–50 mg)	Placebo	No	66.2 (15.3)	30	473 (47%)	Yes	Yes	Yes	Yes
Kline <i>et al</i> ¹⁶	83	Tenecteplase	Placebo	No	55.4 (14)	5	49 (59.0)	Yes	Yes	Yes	Yes
Becattini <i>et al</i> ¹⁷	58	Tenecteplase (30–50 mg)	Placebo	No	68.1 (1.9)	7	13 (22.4)	Yes	Yes	Yes	Yes
Konstantinides <i>et al</i> ¹⁸	256	Alteplase (100 mg)	Placebo	No	62.1 (10.5)	30	122 (47.6)	Yes	Yes	Yes	Yes
Goldhaber <i>et al</i> ¹⁹	101	rt-PA (100 mg)	Placebo	No	58.5 (17)	14	44 (44.0)	Yes	Yes	Yes	Yes
Dalla-Volta <i>et al</i> ²⁰	36	Alteplase (100 mg)	Placebo	No	64.7 (12.5)	30	12 (33.0)	Yes	Yes	Yes	Yes
Levine <i>et al</i> ²¹	58	Alteplase (0.6 mg/kg of ideal body weight)	Placebo	No	61.5 (2.7)	10	29 (54.5)	Yes	Yes	Yes	Yes
PIOPED ²²	13	Alteplase (40–80 mg)	Placebo	No	58.5 (15.8)	7	9 (55.6)	Yes	Yes	Yes	No
Marini <i>et al</i> ²³	30	Urokinase (800 000 IU for 12 hours/day for 3 days or 3 300 000 IU for 12 hours)	Placebo	No	53 (23–72)		11 (44)	Yes	Yes	Yes	Yes
Ly <i>et al</i> ²⁴	25	Streptokinase (250 000 IU loading dose, then 100 000 IU/hours for 72 hours)	Placebo	Yes	53.2 (23–70)	10	11 (44.0)	Yes	Yes	No	No
Tibbitt <i>et al</i> ²⁵	30	Streptokinase (600 000 IU over 30 min through PA catheter followed by 100 000 IU/hours intravenous for 72 hours)	Placebo	Yes	48.7 (25–71)	3	15 (50.0)	Yes	Yes	No	No
UPE ²⁶	160	Urokinase (2000 U/lb, then 2000 U/lb/hour for 12 hours)	Placebo	Yes	†	14	92 (57.3)	Yes	Yes	Yes	Yes
Fasullo <i>et al</i> ²⁷	72	Alteplase (100 mg)	Placebo	No	56.0 (16.1)	180	41 (56.9)	Yes	Yes	Yes	Yes
Jerjes-Sanchez <i>et al</i> ²⁸	8	Streptokinase (1500 000 IU)	Placebo	Yes	51 (22.9)	1–3	5 (63.0)	Yes	Yes	Yes	No
Dotter <i>et al</i> ²⁹	31	Streptokinase (2000 000 to 11 000 000 IU)	Placebo	Yes	Yes	14	‡	Yes	Yes	Yes	Yes
Taherkhani <i>et al</i> ³⁰	50	Alteplase (100 mg) or streptokinase (1500 000 IU)	Placebo	No	55.7 (12.4)	7	20 (40.0)	Yes	Yes	Yes	No
Sharifi <i>et al</i> ³¹	121	rt-PA (50 mg)	Placebo	No	Intervention: 58 ⁹ Control: 59 ¹⁰	840	55 (45.5)	Yes	Yes	Yes	Yes
Kucher <i>et al</i> ³²	59	rt-PA (10–20 mg through PA catheter)	Placebo	No	63 (14)	90	28 (47.5)	Yes	Yes	Yes	Yes
Goldhaber <i>et al</i> ³³	90	rt-PA (100 mg)	rt-PA (0.6 mg/kg with a maximum dose of 50 mg)	Yes	Intervention: 53 ¹⁷ Control: 58 ¹⁶	14	46 (51.1)	Yes	Yes	Yes	Yes
Sors <i>et al</i> ³⁴	53	Alteplase (100 mg)	Alteplase (0.6 mg/kg with a maximum dose of 50 mg)	Yes	Intervention: 69 ¹² Control: 67 ¹⁷	Hospital stay	23 (43.4)	Yes	Yes	Yes	Yes
Wang <i>et al</i> ³⁵	118	rt-PA (100 mg)	rt-PA (50 mg)	Yes	Intervention: 51.9 (13.5) Control: 55.3 (14.1)		69 (58.5)	Yes	Yes	Yes	Yes
Abdelsamad <i>et al</i> ³⁶	40	Streptokinase (1000 000 IU over 1 hour)	Streptokinase (250 000 IU over 30 min, then 100 000 IU/hour over 24 hours)	No	NA		NA	Yes	Yes	No	No

*Patients also received standard anticoagulation.

†Precise ages of patients not provided; 50.6% of patients were younger than 50 years and 49.4% of patients 50 years or older.

‡Unspecified.

ICH, intracranial haemorrhage; IU, international units.; rt-PA, recombinant tissue plasminogen activator; VTE, venous thromboembolism.

Table 2 Summary of direct meta-analysis for all-cause mortality and adverse event outcomes

Intervention	Studies (n)	Active intervention*		Control (placebo unless otherwise noted)*		OR (95% CI)
		No. with event	Total (n)	No. with event	Total (n)	
All-cause mortality						
Full-dose thrombolysis	16	23	1010	42	1006	0.64 (0.37 to 1.09)
Low-dose thrombolysis	1	1	61	3	60	0.32 (0.03 to 3.13)
Catheter-directed thrombolysis	1	0	30	1	29	0.31 (0.01 to 7.96)
Full-dose thrombolysis versus low-dose thrombolysis	4	4	112	7	186	1.04 (0.24 to 4.41)
Major bleeding						
Full-dose thrombolysis	16	99	1010	38	1006	2.39 (1.44 to 3.95)
Low-dose thrombolysis	1	0	61	0	60	Not estimable
Catheter-directed thrombolysis	1	0	30	0	29	Not estimable
Full-dose thrombolysis versus low-dose thrombolysis	4	9	112	7	186	2.26 (0.78 to 6.58)
Intracranial haemorrhage						
Full-dose thrombolysis	14 2	15	983	2	978	3.66 (1.13 to 11.86)
Low-dose thrombolysis	1	0	61	0	60	Not estimable
Catheter-directed thrombolysis	1	0	30	0	29	Not estimable
Full-dose thrombolysis versus low-dose thrombolysis	3	3	97	0	161	6.85 (0.74 to 63.24)
Recurrent VTE						
Full-dose thrombolysis	11	19	945	37	945	0.57 (0.32 to 1.03)
Low-dose thrombolysis	1	0	61	3	60	0.13 (0.01 to 2.64)
Catheter-directed thrombolysis	1	0	30	0	29	Not estimable
Full-dose thrombolysis versus low-dose thrombolysis	3	4	97	6	161	1.35 (0.36 to 5.00)

*Patients also received standard anticoagulation.
VTE, venous thromboembolism.

probability of major bleeding (SUCRA, 0.61), followed by catheter-directed thrombolysis (SUCRA, 0.54) and full-dose thrombolysis (SUCRA, 0.17) (figure 3).

Network meta-analysis: secondary outcomes

In network meta-analysis, compared with anticoagulation, all procedures had 0.48–2.07 odds of being associated with ICH (online supplementary efigure 9). Compared with anticoagulant therapy, low-dose thrombolysis was associated with the lowest

		Odds ratio (95% CI) for dying		
Odds ratio (95% CI) for bleeding	<i>Full-dose thrombolysis</i>	1.28 (0.40 to 4.12)	1.93 (0.07 to 51.33)	0.60 (0.36 to 1.01)
	2.22 (0.71 to 6.89)	<i>Low-dose thrombolysis</i>	1.50 (0.05 to 47.94)	0.47 (0.14 to 1.59)
	2.07 (0.03 to 126.08)	0.93 (0.01 to 65.65)	<i>Catheter-directed thrombolysis</i>	0.31 (0.01 to 7.96)
	2.00 (1.06 to 3.78)	0.90 (0.25 to 3.21)	0.97 (0.02 to 56.03)	<i>Placebo</i>

Figure 2 Network meta-analysis estimates of all-cause mortality (upper triangle) and major bleeding (lower triangle) for each comparison.




Figure 3 Clustered ranking plot based on cluster analysis of SUCRA for benefit (all-cause mortality) and safety (major bleeding). Treatments lying in the upper right corner are more effective and safe than the other treatments. SUCRA, surface under the cumulative ranking curve.

odds of ICH (OR 0.48; 95% CI 0.07 to 3.14; SUCRA, 0.78), whereas full-dose thrombolysis (OR 2.07; 95% CI 0.86 to 5.02; SUCRA, 0.16) was associated with the highest odds of ICH (online supplementary efigure 10).

Compared with anticoagulation, all procedures had 0.34–0.97 lower odds of being associated with recurrent embolism (online supplementary efigure 9). Compared with anticoagulant therapy, low-dose thrombolysis was associated with the lowest odds of recurrent embolism (OR 0.34; 95% CI 0.09 to 1.25; SUCRA, 0.81), whereas catheter-directed thrombolysis (OR 0.97; 95% CI 0.02 to 50.36; SUCRA, 0.40) was associated with the highest odds of recurrent embolism (online supplementary efigure 10).

Sensitivity analysis

Results from sensitivity analyses are reported in online supplementary efigure 1. Overall, the results were similar to the main analysis for the primary outcome in sensitivity analyses based on (1) restricting only to studies in patients with haemodynamically stable PE; (2) restricting only to trials where the mean age of participants in the thrombolytic group was >65 years; and (3) alternative statistical model (frequentist approach using a random-effects inconsistency model).

Publication bias and network consistency

There was no evidence of publication bias, either qualitatively based on funnel-plot asymmetry (online supplementary efigure 11) or quantitatively (Egger regression test, $P > 0.05$ for all comparisons), although the number of studies included in

each comparison was small. There were significant differences between direct and indirect estimates in the only closed loop that allowed assessment of network consistency (anticoagulation–full-dose thrombolysis–low-dose thrombolysis).

Quality of evidence

The risk of bias summary and figure for included studies are listed in online supplementary efigure 12 and efigure 2. Some studies did not present details for randomisation, allocation concealment and blinding. No more than four of the included trials (<20%) were deemed to be at high risk of bias in only three domains (randomisation, allocation concealment, blinding) of the Cochrane Collaboration risk of bias tool. In most domains, the majority of trials were at low risk, except for the allocation concealment and blinding categories in which most trials were at an unclear risk due to inadequate reporting of methods.

DISCUSSION

To our knowledge, this is the first network meta-analysis comparing full-dose thrombolysis, low-dose thrombolysis, catheter-directed thrombolysis and inactive controls on mortality and other adverse outcomes in patients with acute symptomatic PE. The study has several key findings. First, full-dose thrombolysis, low-dose thrombolysis and catheter-directed thrombolysis showed a non-significant trend toward lower risk of all-cause death compared with anticoagulation. Second, full-dose thrombolysis was associated with higher odds of major bleeding compared with anticoagulant treatment, with moderate

confidence in estimates, but was associated with lower odds of recurrences. Third, low-dose thrombolysis was the treatment that performed best in terms of efficacy (all-cause mortality) and safety (major bleeding). However, the clinical interpretation of these findings is limited by the uncertainty around these estimates and by the potential bias due to the small number of trials in each node.

Traditional pairwise meta-analyses are limited in helping to summarise the most effective treatment among different kinds of recanalisation procedures. Other than comparisons between full-dose thrombolysis and anticoagulant therapy,^{37,38} the number of studies that analysed each particular pair of treatments is still relatively small. Furthermore, for some procedures (ie, full-dose vs catheter-directed thrombolysis), there was no direct comparative research. The ability to estimate effectiveness in this work using network meta-analysis allows for more comprehensive assessment of treatment options than has been previously possible. Additionally, in contrast to separate pairwise analyses, we have been able to rank each treatment based on the strength of its association with mortality and bleeding. Even though the results of the pairwise and network meta-analyses were mostly similar, the biggest difference was seen in the comparison of full-dose with anticoagulation on ICH with the pairwise meta-analysis estimating a larger association than the network model. This was most likely because of the large amount of between-study variation observed in the indirect comparisons being incorporated into the analysis.

Some previous pairwise meta-analyses showed significantly lower associated mortality with full-dose systemic thrombolytic use in PE.^{37,38} In our study, we did not find a significant reduction in all-cause mortality with full-dose thrombolytic therapy. This discrepancy between the studies may be explained at least in part by the use of different methodological and statistical techniques. For low-dose systemic thrombolysis and catheter-directed thrombolysis, lack of statistical power might account for the non-significant results, as suggested by the wider CIs. Alternatively, full-dose systemic thrombolysis showed a significant association with major bleeding, a finding consistent with previous meta-analyses.^{37–39} While low-dose and catheter-directed thrombolysis have the potential to offer benefits of full-dose systemic thrombolysis while minimising bleeding risk attributable to a lower dose of the thrombolytic agent, limited randomised clinical trial data might be the main obstacle for providing a definitive conclusion on the comparison of the effect of different reperfusion therapies on major bleeding and ICH. On balance, our results show that low-dose and catheter-directed thrombolysis seem the most highly ranked treatment across the two primary outcomes. Since catheter-directed thrombolysis requires rapid access to the cardiac catheterisation or interventional radiology laboratory,⁵ low-dose thrombolysis is appealing for patients with PE when early recanalisation procedures are indicated. However, it should be kept in mind that, in patients with such presentations, particularly when PE is associated with haemodynamic instability, there are relatively few data for any approaches other than standard-dose systemic thrombolysis.

Ultimately, given the differences in safety, efficacy, and response to therapy, from a clinical perspective, the clinician should always consider the overall clinical picture, and patient management plans need to balance the risks and benefits. There is also a need for randomised trials that compare low-dose thrombolytic therapy with anticoagulation alone in stable patients who have intermediate-risk to high-risk PE. Evidence from such studies would place the role of this procedure for PE on a firmer footing.

This study has limitations. First, there was a paucity of head-to-head trials. Second, the biggest threat to validity of the results of any meta-analysis is conceptual heterogeneity (ie, considerable differences among trials in patient characteristics, studied interventions, outcome assessment or study design), which can limit the comparability of trials. Strategies to limit the effect of conceptual heterogeneity included strict inclusion and exclusion criteria and the use of various sensitivity analyses to assess the robustness of the results. Third, we found inconsistency for efficacy, which was mainly determined by the loop of anticoagulation–full-dose thrombolysis–low-dose thrombolysis. Since some evidence suggests that quality of thrombolytic clinical trials has substantially changed in the past 30 years, we believe that this inconsistency might be a consequence of a cohort effect that relates to different methods used in the older studies compared with those done more recently.⁴⁰ Fourth, ranking probabilities may be affected by unequal numbers of trials per comparison, sample size of individual studies, network configuration and effect sizes among treatments and should be interpreted with caution. Finally, some included trials had an unclear or high rate of selection and performance bias, and there are unaddressed concerns regarding the effect of recanalisation procedures in a clinical setting.

In conclusion, compared with standard anticoagulation, recanalisation procedures had a similar risk of all-cause mortality, though full-dose thrombolysis was associated with an increased risk of major bleeding. This network meta-analysis did not identify a statistically significant difference between the outcomes associated with these therapies, but low-dose thrombolysis was associated with the lowest probability of dying and bleeding. The current body of evidence is limited, and further conclusive studies are needed to establish the role of each of the recanalisation procedures.

Author affiliations

- ¹Respiratory Department, Hospital Ramón y Cajal, Madrid, Spain
²Medicine Department, Universidad de Alcalá (IRYCIS), Alcalá de Henares, Spain
³Faculty of Health Sciences, Universidad Francisco de Vitoria, Pozuelo de Alarcón, Spain
⁴Biostatistics Department, Ramón y Cajal Hospital, IRYCIS, CIBERESP, Madrid, Spain
⁵Department of Thrombosis and Haemostasis, Leiden University Medical Center, Leiden, The Netherlands
⁶Cedars-Sinai Medical Center, Los Angeles, California, USA
⁷Divisions of Pulmonary and Critical Care Medicine and General Medical Sciences, Washington University School of Medicine, St Louis, Missouri, USA

Contributors Study concept and design: DJ, MVH, VT and RDY. Acquisition of data, analysis and interpretation of data and statistical analysis: DJ, CM-S, AM, JZ, RM, DB, FAK, MVH, VT and RDY. Drafting of the manuscript: DJ, FAK, MVH, VT and RDY. Critical revision of the manuscript for important intellectual content: DJ, CM-S, AM, JZ, RM, DB, FAK, MVH, VT and RDY. Study supervision: DJ and RDY. The corresponding author, DJ, had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Funding This study (PII11/00246) was supported by the Instituto de Salud Carlos III (Plan Estatal de I+D+i 2013–2016) and co-financed by the European Development Regional Fund 'A way to achieve Europe'.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Goldhaber SZ, Visani L, De Rosa M. Acute pulmonary embolism: clinical outcomes in the International Cooperative Pulmonary Embolism Registry (ICOPER). *Lancet* 1999;353:1386–9.

- 2 Jiménez D, de Miguel-Díez J, Guijarro R, *et al*. Trends in the management and outcomes of acute pulmonary embolism: analysis from the RIETE registry. *J Am Coll Cardiol* 2016;67:162–70.
- 3 Becattini C, Agnelli G. Risk factors for adverse short-term outcome in patients with pulmonary embolism. *Thromb Res* 2001;103:V239–V244.
- 4 Conget F, Otero R, Jiménez D, *et al*. Short-term clinical outcome after acute symptomatic pulmonary embolism. *Thromb Haemost* 2008;100:937–42.
- 5 Kearon C, Akl EA, Ornelas J, *et al*. Antithrombotic therapy for VTE disease: CHEST guideline expert panel report. *Chest* 2016;149:315–52.
- 6 Meyer G, Vicaut E, Danays T, *et al*. Fibrinolysis for patients with intermediate-risk pulmonary embolism. *N Engl J Med* 2014;370:1402–11.
- 7 Konstantinides SV, Torbicki A, Agnelli G, *et al*. 2014 ESC guidelines on the diagnosis and management of acute pulmonary embolism. *Eur Heart J* 2014;35:3033–73.
- 8 Hutton B, Salanti G, Caldwell DM, *et al*. The PRISMA extension statement for reporting of systematic reviews incorporating network meta-analyses of health care interventions: checklist and explanations. *Ann Intern Med* 2015;162:777–84.
- 9 Lefebvre C, Manheimer E, Glanville J. Chapter 6: Searching for studies. In: Higgins JPT, Green S, eds. *Cochrane handbook for systematic reviews of interventions*. London, England: Cochrane Collaboration, 2011.
- 10 Cochrane Higgins JPT Green S eds. *Cochrane handbook for systematic reviews of interventions* [version 5.1.0, updated March 2011]. <http://handbook.cochrane.org/> (accessed 20 Jun 2016).
- 11 DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7:177–88.
- 12 Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21:1539–58.
- 13 Caldwell DM, Ades AE, Higgins JP. Simultaneous comparison of multiple treatments: combining direct and indirect evidence. *BMJ* 2005;331:897–900.
- 14 White IR, Barrett JK, Jackson D, *et al*. Consistency and inconsistency in network meta-analysis: concepts and models for multi-arm studies. *Res Synth Methods* 2012;3:111–25.
- 15 Salanti G, Ades AE, Ioannidis JP. Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: an overview and tutorial. *J Clin Epidemiol* 2011;64:163–71.
- 16 Kline JA, Nordenholz KE, Courtney DM, *et al*. Treatment of submassive pulmonary embolism with tenecteplase or placebo: cardiopulmonary outcomes at 3 months: multicenter double-blind, placebo-controlled randomized trial. *J Thromb Haemost* 2014;12:459–68.
- 17 Becattini C, Agnelli G, Salvi A, *et al*. Bolus tenecteplase for right ventricle dysfunction in hemodynamically stable patients with pulmonary embolism. *Thromb Res* 2010;125:e82–e86.
- 18 Konstantinides S, Geibel A, Heusel G, *et al*. Heparin plus alteplase compared with heparin alone in patients with submassive pulmonary embolism. *N Engl J Med* 2002;347:1143–50.
- 19 Goldhaber SZ, Haire WD, Feldstein ML, *et al*. Alteplase versus heparin in acute pulmonary embolism: randomised trial assessing right-ventricular function and pulmonary perfusion. *Lancet* 1993;341:507–11.
- 20 Dalla-Volta S, Palla A, Santolicandro A, *et al*. PAIMS 2: alteplase combined with heparin versus heparin in the treatment of acute pulmonary embolism. Plasminogen activator Italian multicenter study 2. *J Am Coll Cardiol* 1992;20:520–6.
- 21 Levine M, Hirsh J, Weitz J, *et al*. A randomized trial of a single bolus dosage regimen of recombinant tissue plasminogen activator in patients with acute pulmonary embolism. *Chest* 1990;98:1473–9.
- 22 Investigators P. Tissue plasminogen activator for the treatment of acute pulmonary embolism: a collaborative study by the PIOPED Investigators. *Chest* 1990;97:528–33.
- 23 Marini C, Di Ricco G, Rossi G, *et al*. Fibrinolytic effects of urokinase and heparin in acute pulmonary embolism: a randomized clinical trial. *Respiration* 1988;54:162–73.
- 24 Ly B, Arnesen H, Eie H, *et al*. A controlled clinical trial of streptokinase and heparin in the treatment of major pulmonary embolism. *Acta Med Scand* 1978;203:465–70.
- 25 Tibbitt DA, Davies JA, Anderson JA, *et al*. Comparison by controlled clinical trial of streptokinase and heparin in treatment of life-threatening pulmonary embolism. *Br Med J* 1974;1:343–7.
- 26 The Urokinase Pulmonary Embolism Trial. Urokinase pulmonary embolism trial. Phase 1 results: a cooperative study. *JAMA* 1970;214:2163–72.
- 27 Fasullo S, Scalzo S, Maringhini G, *et al*. Six-month echocardiographic study in patients with submassive pulmonary embolism and right ventricle dysfunction: comparison of thrombolysis with heparin. *Am J Med Sci* 2011;341:33–9.
- 28 Jerjes-Sanchez C, Ramirez-Rivera A, de Lourdes Garcia M, *et al*. Streptokinase and heparin vs heparin alone in massive pulmonary embolism: a randomized controlled trial. *J Thromb Thrombolysis* 1995;2:227–9.
- 29 Dotter CT, Seaman AJ, Rösch J, *et al*. Streptokinase and heparin in the treatment of pulmonary embolism: a randomized comparison. *Vasc Surg* 1979;13:42–52.
- 30 Taherkhani M, Taherkhani A, Hashemi SR, *et al*. Thrombolytic-plus-anticoagulant therapy versus anticoagulant-alone therapy in submassive pulmonary thromboembolism (TVASPE Study): a randomized clinical trial. *J Tehran Heart Cent* 2014;9:104–8.
- 31 Sharifi M, Bay C, Skrocki L, *et al*. Moderate pulmonary embolism treated with thrombolysis (from the "MOPETT" Trial). *Am J Cardiol* 2013;111:273–7.
- 32 Kucher N, Boekstegers P, Müller OJ, *et al*. Randomized, controlled trial of ultrasound-assisted catheter-directed thrombolysis for acute intermediate-risk pulmonary embolism. *Circulation* 2014;129:479–86.
- 33 Goldhaber SZ, Agnelli G, Levine MN. Reduced dose bolus alteplase vs conventional alteplase infusion for pulmonary embolism thrombolysis. An international multicenter randomized trial. The Bolus Alteplase Pulmonary Embolism Group. *Chest* 1994;106:718–24.
- 34 Sors H, Pacouret G, Azarian R, *et al*. Hemodynamic effects of bolus vs 2-h infusion of alteplase in acute massive pulmonary embolism. A randomized controlled multicenter trial. *Chest* 1994;106:712–7.
- 35 Wang C, Zhai Z, Yang Y, *et al*. Efficacy and safety of low dose recombinant tissue-type plasminogen activator for the treatment of acute pulmonary thromboembolism: a randomized, multicenter, controlled trial. *Chest* 2010;137:254–62.
- 36 Abdelsamad AA, El-Morsi AS, Mansour AE. Efficacy and safety of high dose versus low dose streptokinase for treatment of submassive pulmonary embolism. *Egy Heart J* 2011;63:67–72.
- 37 Chatterjee S, Chakraborty A, Weinberg I, *et al*. Thrombolysis for pulmonary embolism and risk of all-cause mortality, major bleeding, and intracranial hemorrhage: a meta-analysis. *JAMA* 2014;311:2414–21.
- 38 Marti C, John G, Konstantinides S, *et al*. Systemic thrombolytic therapy for acute pulmonary embolism: a systematic review and meta-analysis. *Eur Heart J* 2015;36:605–14.
- 39 Riera-Mestre A, Becattini C, Giustozzi M, *et al*. Thrombolysis in hemodynamically stable patients with acute pulmonary embolism: a meta-analysis. *Thromb Res* 2014;134:1265–71.
- 40 Konstantinides SV. Acute pulmonary embolism revisited. *Heart* 2008;94:795–802.